



Neotropical Helminthology



ORIGINAL ARTICLE / ARTÍCULO ORIGINAL

SEASONAL DYNAMICS OF TWO SPIRURID NEMATODES (NEMATODA: RHABDOCHONIDAE) IN *SCHIZOTHORAX RICHARDSONII* (GRAY, 1832) FROM FIVE COLLECTION SITES OF DISTRICT CHAMBA, HIMACHAL PRADESH, INDIA

DINÁMICA ESTACIONAL DE DOS NEMATODOS ESPIRÚRIDOS (NEMATODA: RHABDOCHONIDAE) EN *SCHIZOTHORAX RICHARDSONII* (GRAY, 1832) DE CINCO SITIOS DE RECOLECCIÓN DEL DISTRITO DE CHAMBA, HIMACHAL PRADESH, INDIA

Suman Kumari^{1*}; Yanchen Dolma¹ & Deepak C. Kalia²

¹ Department of Life-Sciences, MCM DAV College Kangra (H.P.), India 176001.

² Department of Bio-Sciences, Himachal Pradesh University, Summer Hill Shimla (H.P.), India 171005.

* Corresponding author: sumanhpu11@gmail.com

Suman Kumari: <https://orcid.org/0000-0002-3255-8442>

Yanchen Dolma: <https://orcid.org/0000-0001-8862-4181>

Deepak C. Kalia: <https://orcid.org/0000-0002-9478-3899>

ABSTRACT

The present study deals with a seasonal survey of nematode parasites from district Chamba (H.P.) India, and summarizes the data of prevalence, intensity, density and index of infections of *Rhabdochona* (*Rhabdochona*) *denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona* (*Filochona*) *hellichi turkestanica* (Skrjabin, 1917) Moravec, Scholz, Ash and Kar, 2010 in the freshwater fish, *Schizothorax richardsonii* (Gray, 1832). Fish samples were collected from five collection sites of district Chamba *i.e.* (Chamba, (Sukrala, Rari, Saho, and Kheri). During the present study, a total of 293 (92 ♂ + 201 ♀) randomly selected fish hosts were examined, in which 163 fish were infected with two species of nematodes. Seasonal studies for both the species from random-sized hosts exhibited higher prevalence, intensity, density, and index of infection in the summer season and least during the rainy season. Of the two species, *R. (Filochona) hellichi turkestanica* dominated in four localities, with higher values of intensity of infection, density, and index of infection during all three seasons.

Keywords: Nematode – *Rhabdochona* – *Schizothorax* – Seasonal dynamics

Este artículo es publicado por la revista Neotropical Helminthology de la Facultad de Ciencias Naturales y Matemática, Universidad Nacional Federico Villarreal, Lima, Perú auspiciado por la Asociación Peruana de Helminología e Invertebrados Afines (APHIA). Este es un artículo de acceso abierto, distribuido bajo los términos de la licencia Creative Commons Atribución 4.0 Internacional (CC BY 4.0) [<https://creativecommons.org/licenses/by/4.0/deed.es>] que permite el uso, distribución y reproducción en cualquier medio, siempre que la obra original sea debidamente citada de su fuente original.



DOI: <https://dx.doi.org/10.24039/rnh20231711569>

RESUMEN

El presente estudio trata de la evaluación estacional de nematodos parásitos del distrito de Chamba (H.P.) India, y resume los datos de prevalencia, intensidad, densidad e índice de infecciones de *Rhabdochona* (*Rhabdochona*) *denudata* (Dujardin, 1845) Railliet, 1916 y *Rhabdochona* (*Filochona*) *hellichi turkestanica* (Skrjabin, 1917) Moravec, Scholz, Ash & Kar, 2010, en peces de agua dulce, *Schizothorax richardsonii* (Gray, 1832). Se recolectaron muestras de peces de cinco sitios del distrito de Chamba (Chamba, Sukrala, Rari, Saho y Kheri). Durante el presente estudio se examinaron un total de 293 (92 ♂ + 201 ♀) peces huéspedes seleccionados al azar, en los cuales 163 peces estaban infectados con dos especies de nematodos. Los estudios estacionales para ambas especies de hospederos de tamaño aleatorio, exhibieron mayor prevalencia, intensidad, densidad e índice de infección en la temporada de verano y menos durante la temporada de lluvias. De las dos especies, *R. (Filochona) hellichi turkestanica* dominó en cuatro localidades, con mayor valor de intensidad de infección, densidad e índice de infección durante las tres temporadas.

Palabras clave: Nematodo – *Rhabdochona* – *Schizothorax* – Dinámicas estacionales

INTRODUCTION

Fish is a cheap and important source of protein which also contains calcium, lipids, minerals, vitamins and oils with desirably low cholesterol level in the diets of fish lovers (Ugbede *et al.*, 2019). India is third largest producer of fish in the world and second in inland fish production. Fisheries are important for the Indian economy as it provides employment opportunities; is a source of nutritional food and foreign exchanges. The fish is said to be 'gold' from water (Kalse & Bhosle, 2015). These edible fishes are known to harbor a number of helminth parasite which cause deterioration in their health, hence their market and nutritive value is affected. The damage caused by helminthes to their hosts is generally related to intensity of infection and depth of parasite penetration with respect to host tissue. Seasonal fluctuation, locality, age, size and sex of the host also determine the parasitic community diversity and burden (Fartade & Chati, 2016). Humans may contract fish-borne zoonotic nematodes by eating fish that has been undercooked or eaten uncooked. Serious outcomes from human infections are possible, including the unexpected death of infected people (Eiras *et al.*, 2018). *Rhabdochona* spp. in the intestine of snow trout has a long term relationship and is considered as natural infection in cyprinids and zoonotic risk to human (Dujardin, 1845; Railliet & Henry, 1916; Khan *et al.*, 2022).

Nematode cause an economic threat to the market value of fish and infected fillets get rejected and can increase production cost (Abiyu *et al.*, 2020). Nematodes can infect fish as adults but larval stages of nematodes infecting piscivorous birds, mammals or reptiles, or less frequently predatory fish, can also infect fish species.

Ingestion of uncooked infected fish meat poses a zoonotic threat to humans (Leela, 2016).

Schizothorax richardsonii (Gray, 1832), commonly known as snow trout is a hillstream and a high valued fish which is famous and preferred for its excellent taste (Yadav *et al.*, 2014).

Rhabdochona (*Rhabdochona*) *denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona* (*Filochona*) *hellichi turkestanica* (Skrjabin, 1917) Moravec, Scholz, Ash and Kar, 2010 are parasitic species of the genus *Rhabdochona*, a widely diverse group of nematode parasitic in freshwater fish (Skrjabin, 1917; Moravec *et al.*, 2010; Moravec & Adlard, 2016).

The present study examines the prevalence, density/abundance, intensity and index of infection of two species *Rhabdochona* (*Rhabdochona*) *denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona* (*Filochona*) *hellichi turkestanica* (Skrjabin, 1917) Moravec, Scholz, Ash and Kar, 2010 recovered from *S. richardsonii* during three different seasons during 2014.

MATERIALS AND METHODS

During a survey on nematode parasites of *S. richardsonii*, a total of 293 (92 ♂ + 201 ♀) fish hosts were examined from five collection sites (Fig. 1) of district Chamba from Indian hill state of Himachal Pradesh. Chamba district is situated between North latitude 32° 10' and 33° 13' and East longitude 75° 45' and 77° 33'. The principal

river that drains the entire Chamba valley between the Dhauladhar and Pangi hills is the Ravi. It ascends in the mid-Himalaya and Bara Bhanghal ranges. It departs

the neighborhood at Kheri and travels to Shahpur's plains (Himachal Pradesh Development Report, 2014; Balokhra, 2015).




Figure 1. Collection sites.

The fish was purchased from the local fisherman, almost immediate to its capture to avoid the deterioration of health of parasitic fauna. The hosts were dissected along the ventral line. The visceral organs were placed in different containers containing saline solution. The mucosal lining of the intestine was scrapped gently with the help of scalpel whereas, the other visceral organs were thoroughly teased into smaller bits to facilitate the release of the embedded parasites. The specimens were washed thoroughly in saline to remove the debris, if any.

The nematode parasites recovered in saline from the hosts were killed and fixed by pouring boiling hot 3-4% formalin or boiling hot 70% ethyl alcohol on the bulk of specimens retained in the thin film of saline by draining off excess of the saline. This step also caused the almost straightening of the nematodes. The fixed nematodes were preserved in fresh fixative in glass vials and labelled

for date, name, locality of host and location of parasites. Voucher specimens (668) are deposited in the parasitology laboratory, department of Biosciences, Himachal Pradesh University, Shimla.

The nematodes were bulk cleared in lactophenol (Glycerin: 2 parts; Distilled water: 1 part; Lactic acid: 1 part & phenol crystal (melted): 1 part) or glycerin (involving a series of upgrading mixtures of 70% alcohol & glycerin for the preparation of temporary mounts). The parasites were identified following Anderson *et al.* (2009) and Gibbons (2010) for the generic diagnosis of the worms, followed by consultation of the relevant literature available worldwide, for the validation of the species. The data obtained were analyzed following the formula of Tenoza & Zejda (1974), Margolis *et al.* (1982) and Bush *et al.* (1997) to study the following parameters.

$$\text{Prevalence} = \frac{\text{Total number of hosts infected}}{\text{Total number of hosts examined}} \times 100$$

$$\text{Intensity of infection} = \frac{\text{Number of parasites collected}}{\text{Number infected hosts}}$$

$$\text{Density/Abundance} = \frac{\text{Number of parasites collected}}{\text{Number of hosts examined}}$$

Index of infection: It is expressed by the formula of Tenoza & Zejda (1974)

$$Z = \frac{A \times B}{C^2}$$

where, Z = Index of infection

- A = Number of parasites collected
- B = Number of hosts infected
- C = Number of hosts examined

Ethical aspects: The protocol for collection of fishes and nematodes are in accordance with the norms of ethical committee, department of Biosciences, Himachal Pradesh University, Shimla and Himachal Pradesh forest department wildlife wing No. WL/ Research study / 234-36.

RESULTS

During a survey on nematode parasites of *S. richardsonii*, a total of 293 (92 ♂ + 201 ♀) fish hosts were examined from five collection sites of district Chamba: locality I (Chamba), locality II (Sukrala), locality III (Rari), locality IV (Saho) and locality V (Kheri) (Fig. 1) during three (in agreement with Himachal Pradesh Development report, 2014) different seasons; winter season (January-February, 2014), summer season (April-May, 2014) and rainy season (August-September, 2014) with the climate of Himachal Pradesh varying from semi tropical to the semi-arctic depending upon the altitude and the three seasons time period, with the rainy from July to September, winter from October to March and summer from April to June. The recovered nematode parasites (105 ♂ + 599 ♀) were of the two species of *Rhabdochona* Railliet, 1916; *R. (Rhabdochona) denudata* (49 ♂ + 283 ♀) and *R. (Filochona) hellichi turkestanica* (56 ♂ + 316 ♀).

The seasonal dynamics for both the species from randomized hosts, at five different localities exhibited higher prevalence of nematode parasite in summer season (April-May) (55.55%, 61.11%, 84.62%, 86.67%, 40%) (Fig. 2) and the least prevalence during rainy season (August-September) (42.31%, 45.45%, 50%, 55%, 35%), (Fig. 3) with an overall prevalence of 49.25%, 54.76%, 65.95%, 71.83%, 37.87%. The intensity of infection, density or abundance and index of infection for both the species of nematodes were also observed as maximum during summer season with 1.8, 1.82, 4.09, 2.23, 1.16; 1, 1.1, 3.46, 1.93, 0.47; 0.56, 0.68, 2.92, 1.68, 0.19 (Fig.

5, 8, 11) respectively for *R. (Rhabdochona) denudata* and 2.3, 2.45, 2.72, 2.88, 1.83; 1.28, 1.5, 2.31, 2.5, 0.73; 0.71, 0.92, 1.95, 2.17, 0.29 (Fig. 5, 8, 11) respectively for *R. (Filochona) hellichi turkestanica*. The record for rainy season was 1.36, 1.40, 3.33, 1.64, 1.14; 0.58, 0.64, 1.67, 0.9, 0.4; 0.24, 0.29, 0.83, 0.49, 0.14 (Fig. 6, 9, 12) respectively for *R. (Rhabdochona) denudata*, 1916 and 1.82, 2, 2.17, 2.55, 1.29; 0.77, 0.91, 1.08, 1.4, 0.45; 0.33, 0.41, 0.54, 0.77, 0.16 (Fig. 6, 9, 12) respectively for *R. (Filochona) hellichi turkestanica*.

Of the five different localities, the observed prevalence was higher in the locality IV followed by locality III, locality II, locality I and of the least prevalence in locality V, while other parameters, viz., intensity of infection, density/ abundance and index of infection for *R. (R.) denudata* were higher in locality III followed by locality IV, locality II, locality I and locality V and for *R. (F.) hellichi turkestanica* these parameters were higher in locality IV followed by locality III, locality II, locality I and least values were observed in locality V (Tables 1 and 2; Fig. 2-13).

Of the above species, *R. (F.) hellichi turkestanica* dominated in all localities with higher values of intensity of infection (2.28), density or abundance (1.26) and index of infection (0.70), during the three seasons, whereas *R. (R.) denudata* exhibited the value of 2.04, 1.14 and 0.63 for the above said parameters. The infected hosts consisted mainly of females (114) as compared to 49 males, for both the species.

Table 1. Seasonal dynamics of *Rhabdochona (Rhabdochona) demudata* (Dujardin, 1845) Railliet, 1916.

Sr. No.	Localities	Seasons	Months	Number of hosts examined		Number of hosts infected		Number of parasite		Prevalence (%)	Intensity of infection	Density/ Abundance	Index of infection			
				Total	♂	♀	Total	♂	♀					Total	♂	♀
I	Chamba	Summer	April- May	18	6	12	10	3	7	18	3	15	1	0.56		
II	Sukrala	Summer	April- May	18	3	15	11	3	8	20	3	17	1.11	0.68		
III	Rari	Summer	April- May	13	4	9	11	2	9	45	7	38	3.46	2.92		
IV	Saho	Summer	April- May	30	12	18	26	8	18	58	8	50	1.93	1.68		
V	Kheri	Summer	April- May	30	10	20	12	4	8	14	2	12	0.47	0.19		
I	Chamba	Rainy	August-September	26	6	20	11	1	10	15	5	10	0.58	0.24		
II	Sukrala	Rainy	August-September	11	4	7	5	1	4	7	0	7	0.64	0.29		
III	Rari	Rainy	August-September	12	4	8	6	2	4	20	3	17	1.67	0.83		
IV	Saho	Rainy	August-September	20	8	12	11	3	8	18	2	16	0.9	0.49		
V	Kheri	Rainy	August-September	20	5	15	7	2	5	8	0	8	0.4	0.14		
I	Chamba	Winter	January-February	23	9	14	12	4	8	17	3	14	0.74	0.39		
II	Sukrala	Winter	January-February	13	5	8	7	3	4	10	2	8	0.77	0.41		
III	Rari	Winter	January-February	22	6	16	14	6	8	50	5	45	2.27	1.44		
IV	Saho	Winter	January-February	21	6	15	14	5	9	25	5	20	1.19	0.79		
V	Kheri	Winter	January-February	16	4	12	6	2	4	7	1	6	0.44	0.16		
Total				293			163			332			1.14	0.63		
													2.04	0.63		

Table 2. Seasonal dynamics of *Rhabdochona (Filochona) hellichi turkestanica* (Skjabin, 1917) Moravec et al., 2010.

Sr. No.	Localities	Seasons	Months	Number of hosts examined				Number of hosts infected				Prevalence (%)	Intensity of infection	Density/ Abundance	Index of infection	
				Total		♂		Total		♂						
				♂	♀	♂	♀	♂	♀	♂	♀					
I	Chamba	Summer	April- May	18	6	12	10	3	7	23	3	20	55.55	2.3	1.28	0.71
II	Sukrala	Summer	April- May	18	3	15	11	3	8	27	4	23	61.11	2.45	1.5	0.92
III	Rari	Summer	April- May	13	4	9	11	2	9	30	6	24	84.62	2.72	2.31	1.95
IV	Saho	Summer	April- May	30	12	18	26	8	18	75	10	65	86.67	2.88	2.5	2.17
V	Kheri	Summer	April- May	30	10	20	12	4	8	22	3	19	40	1.83	0.73	0.29
I	Chamba	Rainy	August-September	26	6	20	11	1	10	20	3	17	42.31	1.82	0.77	0.33
II	Sukrala	Rainy	August-September	11	4	7	5	1	4	10	0	10	45.45	2	0.91	0.41
III	Rari	Rainy	August-September	12	4	8	6	2	4	13	2	11	50	2.17	1.08	0.54
IV	Saho	Rainy	August-September	20	8	12	11	3	8	28	4	24	55	2.55	1.4	0.77
V	Kheri	Rainy	August-September	20	5	15	7	2	5	9	1	8	35	1.29	0.45	0.16
I	Chamba	Winter	January-February	23	9	14	12	4	8	23	3	20	52.17	1.92	1	0.52
II	Sukrala	Winter	January-February	13	5	8	7	3	4	14	2	12	53.85	2	1.08	0.58
III	Rari	Winter	January-February	22	6	16	14	6	8	30	6	24	63.63	2.14	1.36	0.87
IV	Saho	Winter	January-February	21	6	15	14	5	9	38	7	31	66.66	2.71	1.81	1.21
V	Kheri	Winter	January-February	16	4	12	6	2	4	10	2	8	37.5	1.67	0.63	0.23
Total				293			163			372			55.63	2.28	1.26	0.706




Figure 2. Locality-wise prevalence of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec *et al.*, 2010 in summer season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 3. Locality-wise prevalence of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec *et al.*, 2010 in rainy season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 4. Locality-wise prevalence of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec et al., 2010 in winter season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 5. Locality-wise intensity of infection of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec et al., 2010 in summer season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 6. Locality-wise intensity of infection of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona. (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec *et al.*, 2010 in rainy season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 7. Locality-wise intensity of infection of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona. (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec *et al.*, 2010 in winter season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 8. Locality-wise density/abundance of *Rhabdochona. (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec et al., 2010 in summer season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 9. Locality-wise density/ abundance of *Rhabdochona. (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec et al., 2010 in rainy season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 10. Locality-wise density/ abundance of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona. (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec *et al.*, 2010 in winter season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 11. Locality-wise index of infection of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona. (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec *et al.*, 2010 in summer season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 12. Locality-wise index of infection of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec et al., 2010 in rainy season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 =Rari, Locality 4 = Saho, Locality 5 = Kheri.




Figure 13. Locality-wise index of infection of *Rhabdochona (R.) denudata* (Dujardin, 1845) Railliet, 1916 and *Rhabdochona (F.) hellichi turkestanica* (Skrjabin, 1917) Moravec et al., 2010 in winter season. Locality 1 = Chamba, Locality 2 = Sukrala, Locality 3 = Rari, Locality 4 = Saho, Locality 5 = Kheri.

DISCUSSION

The present result showed that the high prevalence/incidence of infection (86.67%), intensity of infection (4.09, 2.88) and density of infection (3.46, 2.5) for both the species of *Rhabdochona* Railliet, 1916 (Nematoda - Rhabdochonidae) in summer season followed by winter and rainy season, was in agreement with Bhure *et al.* (2010), who while surveying the population ecology of *Rhabdochona* from *Labeo rohita* (Hamilton & Buchanan) indicated that a high incidence of infection (51.78%), intensity of infection (1.18%) and density of infection (0.61%) of *Rhabdochona* spp. occurred in summer season followed by winter season and rainy season. This type of result indicated that environmental factors and feeding habitats are influencing the seasonality of parasitic infection (Deshmukh *et al.*, 2013). Maheboob (2020) made a similar observation for high infections (Intensity, incidence, density and index of infection) of helminth parasite during summer season, followed by winter season and extremely low in monsoon season. It indicated that ecological factors and feeding habitat are affecting the seasonality of parasitic infection either directly or indirectly.

Ezenwaji *et al.* (2005) studied helminth endo-parasites of mochokids in a tropical rainforest river system observed prevalence, mean intensity and abundance of all the endo-parasites to be generally higher in the dry than in the rainy season as the relatively fast flow of water in lotic habitats would inevitably reduce host-parasite contact frequency resulting in low prevalence. Leela & Rao (2014) could observe that *R. garuui* Agrawal, 1965 in *Glossogobius giuris* (Hamilton-Buchanan, 1822) from Lower Monair Dam (Andhra Pradesh) had lowest prevalence (28.92%), mean intensity (1.42) and relative density 0.41 for the premonsoon months, the highest prevalence (46.06%), mean intensity (2.32) and relative density (1.07) for the monsoon months (Agrawal, 1965). Kalse & Bhosale (2015) while studying biodiversity and prevalence of helminth parasites of Girna Dam fishes from February 2010- January 2012, observed that the incidence of nematode infections during 2010-11 was maximum (71.62%) in summer season, followed by (69.33%) in monsoon season and lower (63.51%) in winter season whereas for the year 2011-12 these were 37.5%, 32.60

and 35.05 respectively. Kumar *et al.* (2016) while studying the prevalence of a spirurid nematode *R. (F.) hellichi turkestanica* in *Tor putitora* (Hamilton-Buchanan, 1822) from Seer Khad, district Bilaspur (Himachal Pradesh) during May, August and November 2015, observed a high prevalence (85%) in May and November with mean intensity (4.79) and relative abundance (4.07) being high in the month of November only (Hamilton-Buchanan, 1822).

Hence, the present observations revealed that despite the identical prevalence, because of the sympatric relationship, the mean intensity, density/abundance and index of infection were 1.14-4.09, 0.40-3.46, 0.14-2.92 for *R. (Rhabdochona) denudata* and 1.29-2.88, 0.45-2.5, 0.16-2.17 for *R. (Filochona) hellichi turkestanica*.

ACKNOWLEDGEMENT

The authors are thankful to the University Grant Commission-Basic Scientific Research for financial support.

Author contributions: CRediT (Contributor Roles Taxonomy)

DCK = Deepak C. Kalia
YD = Yanchen Dolma
SK = Suman Kumari

Conceptualization: DCK, YD, SK
Data curation: SK
Formal Analysis: SK
Funding acquisition: SK
Investigation: SK, YD
Methodology: SK, YD
Project administration: DCK, SK
Resources: DCK, YD, SK
Software: SK
Supervision: DCK
Validation: DCK, YD, SK
Visualization: DCK, YD, SK
Writing – original draft: SK
Writing – review & editing: DCK, YD, SK

BIBLIOGRAPHIC REFERENCES

Abiyu, M., Mekonnen, G., & Hailay, K. (2020). Prevalence of intestinal nematode parasites of the Nile Tilapia (*Oreochromis niloticus*) fish species caught from southwestern Part of lake Tana, central Gondar, Ethiopia. *Journal of Aquaculture research & Development*, 11, 1-7.

- Agrawal, V. (1965). Some new nematode parasites from freshwater fishes of Lucknow. *Indian Journal of Helminthology*, 17, 1-17.
- Anderson, R.C., Chabaud, A.G., & Willmott, S. (2009). *Keys to the Nematode Parasites of Vertebrates: Archival Volume*. CABI, Wallingford (UK), pp. 463.
- Balokhra, J.M. (2015). *The wonderland Himachal Pradesh an encyclopedia on a tiny state of Western Himalaya*. HG Publication.
- Bhure, D.B., Nanware, S.S., Kardile, S.P., & Dhondge, R.M. (2010). A survey of the population ecology of *Rhabdochona* Railliet, 1916 (Nematoda- Rhabdochoniidae) from *Labeo rohita* (Ham. And Buch). *The Ecosphere*, 1, 12-14.
- Bush, A.O., Lafferty, K.D., Lotz, J.M., & Shostak, A.W. (1997). Parasitology meets ecology on its own terms: *Margolis et al., Revisited*. *Journal of Parasitology*, 83, 573-583.
- Deshmukh, V.S., Nanware, S.S., Shinde, S.M., & Bhure, D.B. (2013). Studies on prevalence of Cestoda parasites of freshwater fishes from Nanded region (M.S.) India. *Indian Journal of Applied Research*, 3, 1-2.
- Dujardin, F. (1845). *Histoire naturelle des helminthes ouversintestinaux*. Paris xvi+ pp. 654.
- Eiras, J.C., Pavanelli, G.C., Takemoto, R.M., & Nawa, Y. (2018). Fish – borne nematodiasis in South America: neglected emerging diseases. *Journal of Helminthology*, 92, 649-654.
- Ezenwaji, N.E., Aguiwo, J.N., Ilozumba, P.C.O., & Ezenwaji, H.M.G. (2005). Helminth endo-parasites of mochokids in a tropical rainforest river system. *Animal Research International*, 2, 346-352.
- Fartade, A.M., & Chati, R.S. (2016). Population dynamics of helminth parasite in fishes from Solapur and Osmanabad dist. (M.S.) India. *International Journal of Advance Research*, 4, 427-430.
- Gibbons, L.M. (2010). *Keys to the Nematode Parasites of Vertebrates: Supplementary Volume*. Cambridge University Press.
- Hamilton-Buchanan, F. (1822). *An account of the fishes found in the river Ganges and its branches*: viii + 405, 39 pls.
- Himachal Pradesh Development Report. (2014). *State plan division, planning commission, Government of India*.
- Kalse, A., & Bhosale, Y. (2015). Biodiversity and prevalence of Helminth parasites of Girna dam fishes. *International Journal of Life Sciences, Special Issue A3*, 42-48.
- Khan, W., Naqvi, S.M.H.M., Ahmad, N., Kamal, M., Hassan, H., Noor, A., Khan, S., Ahmad, J., Ullah, U., Akhtar, S., & Shadman, M. (2022). Prevalence of rhabdochoniasis in snow trout of river Swat and river Panjkora, Khyber Pakhtunkhwa province, Pakistan. *Brazilian Journal of Biology*, 82, e238874.
- Kumar, R., Thakur, A., & Kalia, D.C. (2016). Prevalence of a spirurid nematode in *Tor putitora* (Hamilton-Buchanan) from Seer Khad, district Bilaspur (Himachal Pradesh). *International Journal of Advanced Science and Research*, 1, 28-30.
- Leela, B., & Rao, K.R. (2014). Nematode parasites in a freshwater fish *Glossogobius giuris* (Hamilton-Buchanan, 1822) at Lower Manair Dam, Karimnagar Dt. Andhra Pradesh, India. *IOSR Journal of Pharmacy and Biological Sciences*, 9, 37-40.
- Leela, B. (2016). Diversity in freshwater fishes order perciformes at lower Manair dam Karimnagar Dt. Telangana state. *International Journal of Innovative Research in Science, Engineering and Technology*, 5, 9306-9315.
- Maheboob, S.I. (2020). Seasonal study of parasitic infection in fresh water fishes from Siddeshwar dam Hingoli District (M. S.) India. *Journal of Emerging Technologies and Innovative Research*, 7, 821-824.
- Margolis, L., Esch, G.W., Holmes, J.C., Kuris, A.M., & Schad, G.A. (1982). The use of ecological terms in parasitology (report of an ad hoc committee of the American Society of Parasitologists). *Journal of Parasitology*, 68, 131-133.
- Moravec, F., & Adlard, R. (2016). Redescription of *Rhabdochona papuensis* (Nematoda: Thelaziodea), a parasite of rainbow fishes (*Melanotaenia* spp.), the first record of the species of *Rhabdochona* in Australia. *Acta Parasitologica*, 61, 820-827.

- Moravec, F., Scholz, T., Ash, A., & Kar, P.K. (2010). New data on the morphology and taxonomy of three species of *Rhabdochona* (Nematoda: Rhabdochonidae) parasitizing fishes in India. *Folia Parasitologica*, 57, 295-306.
- Railliet, A., & Henry, A. (1916). Nouvelles remarques sur les oxyurides. *Comptes Rendus Societe de Biologie*, 79, 247-250.
- Skrjabin, K.I. (1917). [On the recognition of the helminth diseases of some fishes in Russia]. *Arkhiv Veterinar nykh Nauk*, 1-5, 522-543 (in Russia).
- Tenoza, F., & Zejda, J. (1974). The helminth synster of *Clethrionomys glareolus* in a low land forest and its change. *Acta Science Natural Benasque*, 8, 1-48.
- Ugbede, O.V., Oboro, A.O., Martha, H., Awawa, Y.M., Audu, B.A., & Waziri, Z.A. (2019). Medically important parasites of *Clarias garipienus* (catfish) in Nigeria. *Advances in Biotechnology and Microbiology*, 15, 9-14.
- Yadav, C.N.R., Ghosh, T.K., & Subha, B.R. (2014). *General biology and status of Schizothorax richardsonii in Nepal*. *Nepalese Journal of Biosciences*, 4, 34-39.

Received July 31, 2022

Accepted December 26, 2022.