




# Neotropical Helminthology



ORIGINAL ARTICLE / ARTÍCULO ORIGINAL

## GASTROINTESTINAL PARASITISM BY HELMINTHS AND PROTOZOA IN WILD BIRDS RESCUED FROM TRAFFICKING IN RIVERSIDE COMMUNITIES ON THE SÃO FRANCISCO RIVER, SERGIPE, BRAZIL

### PARASITISMO GASTROINTESTINAL POR HELMINTOS E PROTOZOÁRIOS EM AVES SILVESTRES RESGATADAS DO TRÁFICO EM COMUNIDADES RIBEIRINHAS DO RIO SÃO FRANCISCO, SERGIPE, BRASIL

### PARASITISMO GASTROINTESTINAL POR HELMINTOS Y PROTOZOOS EN AVES SILVESTRES RESCATADAS DEL TRÁFICO EN COMUNIDADES RIBERAS DEL RÍO SÃO FRANCISCO, SERGIPE, BRASIL

Anna Luiza Hora dos Santos<sup>1,9</sup>, João Victor Batista dos Santos<sup>1,9</sup>, Sofia Cerqueira Schettino<sup>2</sup>, José Augusto de Santana Júnior<sup>4</sup>, Elpídio Vicente dos Santos Júnior<sup>5</sup>, Lucas de Oliveira Souza<sup>6</sup>, Aline Borba dos Santos<sup>3</sup>, André Mota Alves<sup>7,9</sup> & Victor Fernando Santana Lima<sup>1,7,8,9\*</sup>

<sup>1</sup> Departamento de Medicina Veterinária do Sertão, Universidade Federal de Sergipe, Nossa Senhora da Glória – SE, Brasil.

<sup>2</sup> Departamento de Medicina Veterinária, Universidade Federal de Sergipe, São Cristóvão – SE, Brasil.

<sup>3</sup> Administração Estadual do Meio Ambiente, Governo do Estado de Sergipe, Aracaju – SE, Brasil.

<sup>4</sup> Centro da Terra, Aracaju – SE, Brasil.

<sup>5</sup> Programa de Pós-graduação em Biotecnologia, Universidade Federal de Sergipe, São Cristóvão – SE, Brasil.

<sup>6</sup> Programa de Pós-graduação em Ecologia e Conservação, Universidade Federal de Sergipe, São Cristóvão – SE, Brasil.


<sup>7</sup> Programa de Pós-graduação em Biologia Parasitária, Universidade Federal de Sergipe, São Cristóvão – SE, Brasil.


<sup>8</sup> Programa de Pós-graduação em Ciências Aplicadas à Saúde, Universidade Federal de Sergipe, Lagarto – SE, Brasil.


<sup>9</sup> Laboratório de Doenças Parasitárias dos Animais, Universidade Federal de Sergipe, Nossa Senhora da Glória – SE, Brasil.

\* Corresponding author: victor.fslima@gmail.com


Anna Luiza Hora dos Santos:  <https://orcid.org/0000-0002-4031-5045>


João Victor Batista dos Santos:  <https://orcid.org/0000-0002-6554-3597>


Sofia Cerqueira Schettino:  <https://orcid.org/0009-0003-5869-1262>


José Augusto de Santana Júnior:  <https://orcid.org/0000-0003-2817-0370>

Elpídio Vicente dos Santos Júnior:  <https://orcid.org/0009-0003-5869-1262>

Lucas de Oliveira Souza:  <https://orcid.org/0000-0001-8917-6069>

Aline Borba dos Santos:  <https://orcid.org/0000-0002-5472-6822>

André Mota Alves:  <https://orcid.org/0000-0002-9150-5287>

Victor Fernando Santana Lima:  <https://orcid.org/0000-0002-7255-0664>

Este artículo es publicado por la revista Neotropical Helminthology de la Facultad de Ciencias Naturales y Matemática, Universidad Nacional Federico Villarreal, Lima, Perú auspiciado por la Asociación Peruana de Helmintología e Invertebrados Afines (APHIA). Este es un artículo de acceso abierto, distribuido bajo los términos de la licencia Creative Commons Atribución 4.0 Internacional (CC BY 4.0) [<https://creativecommons.org/licenses/by/4.0/deed.es>] que permite el uso, distribución y reproducción en cualquier medio, siempre que la obra original sea debidamente citada de su fuente original.



DOI: <https://dx.doi.org/10.62429/rnh20242181784>

## ABSTRACT

Illegal wildlife trafficking is a practice that represents a significant threat to biodiversity and public health in many regions of the world. Birds that are victims of trafficking are subjected to conditions of stress and confinement that favor the proliferation of parasites, compromising animal health and well-being. Furthermore, some of these parasites have zoonotic potential, posing an additional risk to human health. Therefore, this study aims to identify the gastrointestinal parasites present in wild birds originating from trafficking in the São Francisco River Basin region, providing important information about the risks to bird health and human health. Fecal samples were collected from wild birds (n=80) from the orders Passeriformes, Columbiformes, and Psittaciformes. The samples were analyzed using two parasitological techniques: Mini – FLOTAC® and centrifugal-sedimentation with Ziehl-Neelsen staining, to identify the presence of gastrointestinal parasites. The results indicate a prevalence of helminths, such as *Capillaria* sp. (1.5%) and Trichostrongylidae (32.8%), and protozoa, such as *Eimeria* sp. (5%), *Isoospora* sp. (53.7%), *Cryptosporidium* sp. (83.7) and *Entamoeba* sp. (44.8%). Furthermore, these parasites compromise animal health and well-being, causing clinical signs such as diarrhea, loss of appetite, weight loss, and death. In short, identifying gastrointestinal parasites in wild birds rescued from trafficking in riverside communities is essential to understanding the risks to bird health and public health in the region.

**Keywords:** birdlife – helminths – protozoa – zoonosis

## RESUMO

O tráfico ilegal de animais silvestres é uma prática que representa uma ameaça significativa para a biodiversidade e para a saúde pública em muitas regiões do mundo. As aves vítimas do tráfico são submetidas a condições de estresse e confinamento que favorecem a proliferação dos parasitos, comprometendo a saúde e o bem-estar animal. Além disso, alguns desses parasitos apresentam potencial zoonótico, sendo um risco adicional para a saúde humana. Diante de exposto, o objetivo desse estudo é relatar o parasitismo gastrointestinal por helmintos e protozoários em aves silvestres resgatadas do tráfico de animais em comunidades ribeirinhas do Rio São Francisco, Sergipe, Brasil, fornecendo importantes informações sobre os riscos à saúde das aves e à saúde humana. Foram coletadas amostras fecais de aves silvestres (n=80) pertencentes às ordens Passeriformes, Columbiformes e Psittaciformes. As amostras foram analisadas a partir de duas técnicas parasitológicas: Mini – FLOTAC® e centrífugo-sedimentação com coloração de Ziehl-Neelsen, para identificar a presença de parasitos gastrointestinais. Os resultados indicam parasitismo por helmintos, como *Capillaria* sp. (1.5%) e Trichostrongylidae (32.8%), e protozoários, como *Eimeria* sp. (5%), *Isoospora* sp. (53,7%), *Cryptosporidium* sp. (83,7) e *Entamoeba* sp. (44,8%). Esses parasitos comprometem a saúde e bem-estar animal, ocasionando sinais clínicos como diarréia, inapetência, perda de peso e morte. Em suma, a identificação de parasitos gastrointestinais em aves silvestres resgatadas do tráfico em comunidades ribeirinhas é essencial para compreender os riscos à saúde das aves e à saúde pública na região. Além disso, destaca a necessidade de medidas de controle e prevenção das parasitoses, visando promover a conservação da biodiversidade e a sustentabilidade dos ecossistemas ribeirinhos.

**Palavras-chave:** avifauna – helmintos – protozoários – zoonoses

## RESUMEN

El tráfico ilegal de animales silvestres es una práctica que representa una amenaza significativa para la biodiversidad y la salud pública en muchas regiones del mundo. Las aves víctimas del tráfico son sometidas a condiciones de estrés y confinamiento que favorecen la proliferación de parásitos, comprometiendo la salud y el bienestar animal. Además, algunos de estos parásitos presentan potencial zoonótico, siendo un riesgo adicional para la salud humana. Ante lo expuesto, el objetivo de este estudio es reportar el parasitismo gastrointestinal por helmintos y protozoarios en aves silvestres rescatadas del tráfico de animales en comunidades ribereñas del Río São Francisco, Sergipe, Brasil, proporcionando información importante sobre los riesgos para la salud de las aves y la salud humana. Se recolectaron muestras fecales de aves silvestres (n=80) pertenecientes a las órdenes Passeriformes, Columbiformes y Psittaciformes. Las muestras fueron analizadas mediante dos técnicas parasitológicas: Mini-FLOTAC® y centrifugación-sedimentación con coloración de Ziehl-Neelsen, para identificar la presencia de parásitos gastrointestinales. Los resultados indican parasitismo por helmintos, como *Capillaria* sp. (1,5%) y Trichostrongylidae (32,8%), y protozoarios, como *Eimeria* sp. (5%), *Isoospora* sp. (53,7%), *Cryptosporidium* sp. (83,7) y *Entamoeba* sp. (44,8%). Asimismo, estos parásitos comprometen la salud y

el bienestar animal, ocasionando signos clínicos como diarrea, inapetencia, pérdida de peso y muerte. En resumen, la identificación de parásitos gastrointestinales en aves silvestres rescatadas del tráfico en comunidades ribereñas es esencial para comprender los riesgos para la salud de las aves y la salud pública en la región. Además, destaca la necesidad de medidas de control y prevención de parasitosis, con el fin de promover la conservación de la biodiversidad y la sostenibilidad de los ecosistemas ribereños.

**Palabras clave:** avifauna – helmintos – protozoarios – zoonosis

## INTRODUCTION

Brazil stands out as a country that is home to a significant portion of global biodiversity, due to the vast expanses that comprise different biomes (Charity & Ferreira, 2020). However, among the persistent threats to this biodiversity, wildlife trafficking stands out (ICMBio, 2018). The increase in surveillance regarding this practice in Brazil reveals a notable increase in seizures made by environmental agencies, concomitantly highlighting the limitations in effectively combating this illegal activity (Rabelo *et al.*, 2015; Miranda & Dos Anjos, 2022).

In this context, birds stand out as the main targets of wildlife trafficking, due to their notable diversity of species, variation in plumage colors, and richness of songs, combined with their wide geographic distribution (Ribeiro & Silva, 2007; RENCITAS, 2014). After being seized and undergoing rehabilitation processes, the success of their return to their natural habitat is linked to several factors, such as the expression of natural behaviors, the genetic characteristics of the rescued species, and the maintenance of health (De Azevedo *et al.*, 2016).

The susceptibility of wild birds to parasitism is influenced by several factors, including their species, contacts, and the conditions of the environment in which they live. Furthermore, birds face significant challenges in captivity, where stress becomes an additional catalyst for the development of parasites (Boll *et al.*, 2017).

Gastrointestinal parasites are considerable threats, resulting in high rates of morbidity and mortality. Clinical signs associated with these infestations include diarrhea, loss of appetite, weight loss, and anemia, contributing to a negative impact on the health and well-being of birds and, consequently, on the conservation of the species (Reed *et al.*, 2012). Furthermore, some of these etiological agents have a zoonotic potential and can also compromise public health, generating social, economic, and environmental impacts (Saegerman *et al.*, 2012).

For laboratory diagnosis, fecal parasitological analysis methods are used, with the Mini – FLOTAC® technique being a prominent option. This technique makes it possible to identify parasite eggs, oocysts, and larvae, and also determine the parasite load (Cringoli *et al.*, 2017). Another method used is centrifugal sedimentation with Ziehl-Neelsen staining, which can be used to diagnose infections by the potentially zoonotic protozoan *Cryptosporidium* sp. (Silva *et al.*, 2016).

Considering the damage caused by gastrointestinal parasites, carrying out new studies becomes essential to contribute to the preservation of threatened species. Given the above, the present study aims to identify gastrointestinal parasites in wild birds rescued from trafficking in riverside communities along the São Francisco River, Sergipe.

## MATERIAL AND METHODS

In this work, a retrospective study was carried out on the medical records of wild birds victims of trafficking referred and treated by the multidisciplinary team of the 7th stage of the Integrated Preventive Inspection (FPI) in the state of Sergipe (10°54'40" S and 37°04'18" W), Northeast of Brazil in the year 2023. FPI actions took place in 14 municipalities in Sergipe in the São Francisco River Basin: Amparo de São Francisco (10° 9' 58" S and 36° 56' 5" W), Aquidabá (10° 16' 58" S and 37° 1' 12" W), Capela (10° 30' 38" S and 37° 3' 18" W), Graccho Cardoso (10° 13' 42" S and 37° 12' 9" W), Japarutuba (10° 35' 43" S and 36° 56' 24" W), Malhada dos Bois (10° 20' 53" S and 36° 55' 23" W), Pirambu (10° 40' 42" S and 36° 52' 25" W), Nossa Senhora da Glória (10° 13' 0" S and 37° 25' 27" W), Monte Alegre (10° 1' 35" S and 37° 33' 31" W), Nossa Senhora de Lourdes (10° 19' 12" S and 36° 34' 44" W), Itabi (10° 7' 11" S and 37° 6' 18" W), Canhoba (10° 8' 15" and 36° 59' 16" W), Feira

Nova (10° 16' 4" S and 37° 18' 37" W ) and Muribeca (10° 25' 39" S and 36° 57' 45" W).

The experimental design used in this study is classified as cross-sectional with a collection of fecal samples from birds that were victims of trafficking and identification of gastrointestinal parasites present at a single moment in municipalities visited during the 7<sup>th</sup> stage of FPI in the state of Sergipe.

Pools of fecal samples (n=80) were collected through spontaneous defecation from 400 wild birds (400/80 = 5 bird/pool), rescued during actions carried out by

the Fauna Team in Integrated Preventive Inspection (FPI), during the year 2023, in the state of Sergipe. The specimens belonged to the following three orders: Columbiformes (n = 20), Psittaciformes (n = 50), and Passeriformes (n = 330) (Table 1) (Fig. 1). The *pools* were obtained from specimens of different species, ages and sexes, depending on the number of individuals present in the enclosure. Fecal samples were collected through spontaneous defecation, with only the fecal material present on the floor of the cages being collected, after clinical evaluation of the animals. All samples were identified, placed in collection tubes, and refrigerated at 8°C until processing.

**Table 1:** Number of bird species and wild animals included in the study.

Order	Species	Number of specimens
Columbiformes	<i>Columbina squammata</i> (Lesson, 1831)	20
Passeriformes	<i>Cacicus cela</i> (Linnaeus, 1758), <i>Coryphospingus pileatus</i> (Wied, 1821), <i>Cyanoloxia brissonii</i> (Lichtenstein, 1823), <i>Dolospingus fringilloides</i> (Pelzeln, 1870), <i>Euphonia chlorotica</i> (Linnaeus, 1766), <i>Gnorimopsar chopi</i> (Vieillot, 1819), <i>Icterus jamacaii</i> (Gmelin, 1788), <i>Icterus pyrrhopterus</i> (Vieillot, 1819), <i>Mimus gilvus</i> (Vieillot, 1807), <i>Paroaria coronata</i> (Miller, 1776), <i>Piranga flava</i> (Vieillot, 1822), <i>Schistochlamys ruficapillus</i> (Vieillot, 1817) Genus Sicalis, <i>Sicalis flaveola</i> (Linnaeus, 1766), <i>Sicalis luteola</i> (Sparman, 1789), <i>Spinus magellanica</i> (Vieillot, 1805), Genus Sporophila, <i>Sporophila albogularis</i> (Spix, 1825), <i>Sporophila angolensis</i> (Linnaeus, 1766), <i>Sporophila bouvreuil</i> (Muller, 1776), <i>Sporophila lineola</i> (Linnaeus, 1758), <i>Sporophila nigricolis</i> (Vieillot, 1823) , <i>Thraupis sayaca</i> (Linnaeus, 1766), <i>Turdus rufiventris</i> (Linnaeus, 1766), <i>Volatinia jacarina</i> (Linnaeus, 1766) and <i>Zonotrichia capensis</i> (Muller, 1776)	330
Psittaciformes	<i>Alipiopsitta xanthops</i> (Spix, 1824), <i>Amazona amazonica</i> (Linnaeus, 1766), <i>Aratinga auricapillus</i> (Kuhl, 1820), <i>Eupsittula aurea</i> (Gmelin, 1788) and <i>Primolius maracana</i> (Vieillot, 1816)	50



**Figure 1.** Wild birds of the Psittaciformes and Columbiformes orders rescued from animal trafficking in riverside communities on the São Francisco River, Sergipe, Brazil (A) Birds of the species *Eupsittula aurea* (B) Specimens of the species *Columbina squammata*.

To compile the data, the following information was obtained, analyzed and grouped: general taxonomic data of the bird, requested examination, parasitological technique performed, structure and gastrointestinal parasite detected (Henriksen & Pohlenz, 1981; Cringoli *et al.*, 2013). Records with incomplete data were classified as "not noteworthy" (NWN) and were not included in this study. All statistical data were analyzed by InStat GraphPad software (GraphPad Software, 2000), with a significance level of  $p < 0.05$ .


#### Ethical Aspects

This study was carried out with ethical aspects since secondary data from an action by the Public Ministry of Sergipe (MP) were used, which authorized the analysis and use of the data.

#### RESULTS

Overall, 83.7% (67/80) of the samples analyzed were positive for gastrointestinal parasites, regardless of the diagnostic technique. Monoparasitism was observed in 59.7% ( $n=40$ ), while co-infections were identified in 40.3% ( $n=27$ ) of the animals. Among the positive samples, 4.5% were in birds of the order Columbiformes, 10.4% in Psittaciformes and 85.1% in Passeriformes.

In the birds rescued during the inspection action, six types of gastrointestinal endoparasites were identified, belonging to the phyla Nematoda and Protozoa. Protozoa were the most frequent, with emphasis on oocysts of *Isospora* sp. (53.7%), *Eimeria* sp. (5%), *Cryptosporidium* sp. (83.7%), in addition to *Entamoeba* sp. (44.8%). Regarding the phylum Nematoda, nematode eggs from the family Trichostrongylidae (32.8%) and the genus *Capillaria* (1.5%) were detected in the samples (Fig. 2).



**Figure 2.** Egg and oocysts of gastrointestinal parasites identified in fecal samples from birds rescued from trafficking in riverside communities on the São Francisco River (A) Nematode egg from the Trichostrongylidae Family and (B) Oocysts of the protozoan *Cryptosporidium* sp.

In passerine birds, the following gastrointestinal endoparasites were diagnosed: oocysts of *Isoospora* sp. (65.1%), *Cryptosporidium* sp. (86.36%), *Entamoeba* sp. (39.4%) and Trichostrongylidae eggs (28.8%). In Columbiformes birds (*Columbina squammata*), two types of gastrointestinal parasites were identified: *Eimeria* sp. (75%) and *Cryptosporidium* sp. (25%). Finally, five types of enteroparasites were detected in Psittaciformes: *Capillaria* sp. (5.9%), *Eimeria* sp. (36.8%), *Cryptosporidium* sp. (63.2%), *Entamoeba* sp. (57.3%) and Trichostrongylidae (19.1%).

Regarding the clinical aspects of birds infected and/or parasitized by helminths and protozoa, different clinical signs were observed, with emphasis on apathy, diarrhea, inappetence, cachexia, prostration, and ruffled feathers.

## DISCUSSION

Parasitic diseases represent one of the most common and significant health issues affecting wild birds. In captivity, they are exposed to handling conditions that make them susceptible to gastrointestinal parasites (Boll *et al.*, 2017). These conditions include stress, overcrowding, poor hygiene, malnutrition, and space constraints (Sprenger *et al.*, 2018).

Furthermore, it is important to highlight that gastrointestinal parasites can occur asymptotically. However, when there is a high parasite load, affected animals show signs such as apathy, diarrhea, ruffled feathers, prostration, loss of appetite, and cachexia (Cubas *et al.*, 2014). These changes were observed in birds evaluated in this study. Other clinical signs resulting from infection with gastrointestinal parasites include intussusception, intestinal obstruction, and, in more severe cases, death (Papini *et al.*, 2012; Lima *et al.*, 2017).

Regarding the protozoan *Cryptosporidium* sp., a positivity of 83.75% (67/80) was identified in samples from the three orders studied. Passeriformes presented a positivity rate equal to 86.36% (57/66), a higher result than that of Psittaciformes, whose positivity was 63.23% (8/9), and Columbiformes, which was equal to 25% (1/4). Cryptosporidiosis has been reported in several species of wild and captive birds around the world (Ryan, 2010). In general, the infection is asymptomatic, but in immunosuppressed individuals, it causes enterocolitis, diarrhea, and death. Considering that cryptosporidiosis is a zoonotic disease, it is necessary to adopt preventive measures to avoid human and animal infection (Snak *et al.*, 2015).

The coccidia *Isoospora* sp. was detected exclusively in the fecal samples of the Passeriformes evaluated, with positivity equal to 65.1% (43/66). According to Knight *et al.* (2018), *Isoospora* sp. is the coccidia that has the highest prevalence in birds of the order Passeriformes, which reinforces the data obtained in this study. The clinical manifestations of isosporiasis in Passeriformes include weight loss, malabsorption, poor digestion, and even death, signs observed in the birds treated (Giacomo *et al.*, 1997).

Another coccidia, *Eimeria* sp., was identified in Columbiformes and Psittaciformes birds, with positivity equal to 75.0% (3/4) and 11.1% (1/9), respectively. This genus of protozoan is commonly found in Columbiformes, including the species evaluated in this study, *Columbina talpacoti* (Jamriska & Modry, 2012). It is worth highlighting that, in general, coccidiosis causes clinical signs such as diarrhea, apathy, progressive weight loss, ruffled feathers, lethargy, feces stuck to the cloaca, dehydration, weakness, and death, especially in young birds, manifestations commonly observed in the animals evaluated. In this study (Marietto – Gonçalves *et al.*, 2009; Barreto, 2014).

The protozoan *Entamoeba* sp. was identified in the orders Passeriformes and Psittaciformes, with positivity of 39.4% (26/66) and 57.3% (4/9), respectively. Studies that identified gastrointestinal parasites in the same orders of birds corroborate these findings (Sousa *et al.*, 2018; Souza *et al.*, 2019; Prazeres Júnior *et al.*, 2024). According to Graczyk *et al.* (2008), the high frequency of *Entamoeba* sp. in birds, indicates a high degree of anthropization in the environment. Therefore, the presence of this protozoan in fecal samples from birds that are victims of trafficking constitutes yet another problem with this illegal activity.

The nematode *Capillaria* sp. was identified in fecal samples from a specimen of *Amazona amazonica* (5.8%). Santos *et al.* (2022) described the presence of the parasite in this species in a study also carried out in the state of Sergipe. The clinical signs of capillariasis are hemorrhagic diarrhea, anorexia, loss of appetite, progressive weight loss, and death (Gómez *et al.*, 1993).

Trichostrongylidae eggs were identified in the Passeriformes and Psittaciformes samples, with positivity of 28.79% (19/66) and 19.11% (2/9), respectively. The data corroborate the study carried out by Lima *et al.* (2017), who identified this parasite in fecal samples from captive birds in the state of Sergipe. Trichostrongylidae is one of the most common gastrointestinal parasites in birds living in captivity, mainly immuno-

suppressed Columbiformes and Psittaciformes (Freitas, 2002).

In the present study, the centrifugal-sedimentation method with Ziehl – Neelsen staining showed a higher level of positivity (16.2%) compared to the Mini – FLOTAC© technique (12.5%). The centrifuge-sedimentation method is easy to perform, quick, low cost, and has good sensitivity, being the most used method in Brazil for the diagnosis of cryptosporidiosis (Ortolani, 2000).

Wild animals can be carriers and reservoirs of parasitic diseases, affecting the health of natural and domestic ecosystems (Freitas, 2002). Some gastrointestinal parasites identified in the seized birds evaluated in this study have zoonotic potential, such as *Cryptosporidium* sp., which in immunosuppressed individuals can cause death. It is crucial to carry out parasitological studies to identify and control parasites that can affect birds, preventing their spread to other species, including humans (Papini *et al.*, 2012).

Therefore, the identification of gastrointestinal parasites in wild birds rescued from riverside communities on the São Francisco River in the state of Sergipe, Brazil is relevant to understanding the impacts of illegal bird trafficking on the health of poultry populations and the public health of these communities. By identifying the parasites present in birds, especially those with zoonotic potential, such as nematodes and protozoa, the risks to human health can be assessed. Furthermore, the study can contribute to the development of conservation and environmental education strategies in riverside regions, protecting local biodiversity and promoting awareness about the negative impacts of wildlife trafficking.

## ACKNOWLEDGMENTS

We thank the general coordination of Integrated Preventive Inspection: São Francisco River Basin Committee, Federal Public Ministry, State Public Ministry of Sergipe,

and Public Ministry of Labor; to Mr. Antônio Luiz dos Santos and Mr. Jailson Nunes dos Santos; to the Federal Highway Police and all collaborators of the Fauna Team; and to other bodies and entities involved in holding FPI/SE 2023.

## Author contributions: CRediT (Contributor Roles Taxonomy)

**ALHS** = Anna Luiza Hora dos Santos

**JVBS** = João Victor Batista dos Santos

**SCS** = Sofia Cerqueira Schettino

**JASJ** = José Augusto de Santana Júnior

**EVSJ** = Elpídio Vicente dos Santos Júnior

**LOS** = Lucas de Oliveira Souza

**ABS** = Aline Borba dos Santos

**AMA** = André Mota Alves

**VFSL** = Victor Fernando Santana Lima

**Conceptualization:** VFSL, JVBS, ALHS, SCS

**Data curation:** VFSL, JVBS, ALHS, SCS

**Formal Analysis:** VFSL, JVBS, ALHS, SCS

**Funding acquisition:** ALHS, JVBS, JASJ, EVSJ, LOS, ABS, AMA

**Investigation:** VFSL, JVBS, ALHS, SCS

**Methodology:** VFSL, JVBS, ALHS, SCS

**Project administration:** ALHS, JVBS, JASJ, EVSJ, LOS, ABS, AMA

**Resources:** ALHS, JVBS, JASJ, EVSJ, LOS, ABS, AMA

**Software:** ALHS, JVBS, JASJ, EVSJ, LOS, ABS, AMA

**Supervision:** VFSL

**Validation:** VFSL

**Visualization:** VFSL, AMA

**Writing – original draft:** ALHS, JVBS, JASJ, EVSJ, LOS, ABS, AMA

**Writing – review & editing:** ALHS, JVBS, JASJ, EVSJ, LOS, ABS, AMA

## BIBLIOGRAPHIC REFERENCES

Barreto, C., Vilela, D.A.R., Houri, B.F., Lara, L.B., Torres, A.C.D., Silva, A.S.G., Castro Filho, R.P.L., Costa, C.S., & Martins, N.D.S. (2020). New *Isospora* and host species in Brazilian passerines. *Brazilian Journal of Poultry Science*, 22, eRBCA-2019.

- Boll, A.S., Marques, S.M.T., & Alievi, M.M. (2017). Parasitas em Passeriformes e Psittaciformes alojados em centro de triagem no Zoológico de Sapucaia do Sul, Rio Grande do Sul, Brasil. *Revista Portuguesa de Ciências Veterinárias*, 112, 28-34.
- Charity, S., & Ferreira, J.M. (2020). *Wildlife Trafficking in Brazil*. TRAFFIC International, 140. [https://www.traffic.org/site/assets/files/13031/brazil\\_wildlife\\_trafficking\\_assessment.pdf](https://www.traffic.org/site/assets/files/13031/brazil_wildlife_trafficking_assessment.pdf)
- Cringoli, G., Rinaldi, L., Albonico, M., Bergquist, R., & Utzinger, J. (2013). Geospatial (s) tools: integration of advanced epidemiological sampling and novel diagnostics. *Geospatial health*, 7, 399-404.
- Cringoli, G., Maurelli, M.P., Levecke, B., Bosco, A., Vercruysse, J., Utzinger, J., & Rinaldi, L. (2017). The Mini-FLOTAC technique for the diagnosis of helminth and protozoan infections in humans and animals. *Nature protocols*, 12, 1723-1732.
- Cubas, Z.S., Silva, J.C.R., & Catão-Dias, J.L. (2014). *Tratado de Animais Selvagens - Medicina Veterinária*, vol.1. Editora Roca, São Paulo.
- De Azevedo, A.S., Da Silva, G.P., & Braga, G.M.S. (2016). Manejo de fauna apreendida no município de Imperatriz, região sudoeste, do estado do Maranhão. *Pubvet*, 11, 1098-1103.
- Freitas, M.F.L. (2002). Parasitos gastrointestinais de aves silvestres em cativeiro em el estado de Pernambuco, Brasil. *Parasitología Latinoamericana*, 57, 50-54.
- Giacomo, R. (1997). Mortality in black siskins (*Carduelis atrata*) with systemic coccidiosis. *Journal of Wildlife Diseases*, 33, 152-57.
- Gómez M.P.I., Rodríguez-Osorio M., & Aranda-Maza, F. (1993). Parasitacion of falconiform, strigiform and passeriform (Corvidae) birds by helminths in Spain. *Research and Reviews in Parasitology*, 53, 129-135.
- Graczyk, T.K., Majewska, A.C., & Schwab, K.J. (2008). The role of birds in dissemination of human waterborne enteropathogens. *Trends in Parasitology*, 24, 55-59.
- GraphPad Software. (2000). GraphPad Instat software. <https://www.graphpad.com/scientific-software/instat/>
- Henriksen, S.A., & Pohlenz, J.F.L. (1981). Staining of cryptosporidia by a modified Ziehl-Neelsen technique. *Acta Veterinaria Scandinavica*, 22, 594.
- ICMBio. (2018). PRIM – Plano de Redução de Impactos à Biodiversidade, Brasil, consultado em 28 de agosto de 2023, [/www.icmbio.gov.br/portal/faunabrasileira/planos-de-reducao-de-impact](http://www.icmbio.gov.br/portal/faunabrasileira/planos-de-reducao-de-impact).
- Jamriška, J., & Modrý, D. (2012). A new species of *Eimeria* Schneider, 1875 (Apicomplexa: Eimeriidae) from the common wood pigeon *Columba palumbus* Linnaeus, 1758 (Aves: Columbidae). *Acta Protozoologica*, 51, 329-333.
- Knight, A., Ewen J.G., Brekke P., & Santure A.W. (2018). The evolutionary biology, ecology and epidemiology of coccidia of passerine birds. *Advances in Parasitology*, 99, 35-60.
- Lima, V.F.S., Cringoli, G., Rinaldi, L., Monteiro, M.F.M., Calado, A.M.C., Ramos, R.A.N., Meira-Santos, P.O., & Alves, L.C. (2015). A comparison of mini-FLOTAC and FLOTAC with classic methods to diagnosing intestinal parasites of dogs from Brazil. *Parasitology research*, 114, 3529-3533.
- Marietto-Gonçalves, G.A., Martins, T.F., de Lima, E.T., Lopes, R.S., & Andreatti Filho, R.L. (2009). Prevalência de endoparasitas em amostras fecais de aves silvestres e exóticas examinadas no Laboratório de Ornitopatologia e no Laboratório de Enfermidades Parasitárias da FMVZ-UNESP/Botucatu-SP. *Brazilian Animal Science*, 10, 349-354.
- Miranda, D.F., & Anjos, J. (2022). Análise Da Lista Vermelha De Espécies De Aves Da Caatinga, Brasil. *Revista Multidisciplinar de Educação e Meio Ambiente*, 3, 1-9.



- Ortolani, E. L. (2000). Standardization of the modified ZiehlNeelsen technique to stain oocysts of *Cryptosporidium* sp. *Revista Brasileira de Parasitologia Veterinária*, 9, 29-31.
- Papini, R., Girivetto, M., Marangi, M., Mancianti, F., & Giangaspero, A. (2012). Endoparasite infections in pet and zoo birds in Italy. *The Scientific World Journal*, 1-9. <https://doi.org/10.1100/2012/253127>
- Prazeres Júnior, F.R., Alves, A.M., Oliveira Neto, M.B., Oliveira, M.R., Da Silva, W.S.I., De Jesus, J.V., Barata, C.V., Silva, L.M.S., Gomes, A.C., Moreira, A.C., Pereira, L.M.F., & Lima, V.F.S. (2024). Survey on helminths and protozoa of captive wild and exotic birds from Northeastern Brazil. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 76, e13083.
- Rabelo, F.R., De Oliveira, A.G., & Machado, R.A.S. (2015). Tipologia dos crimes ambientais referentes à fauna silvestre no estado da Bahia: uma análise dos autos de infração do INEMA e do IBAMA entre 2001 e 2015. *Sitientibus*, 53, 18-22.
- Reed, T.E., Daunt, F., Kiploks, A.J., Burthe, S.J., Granroth-Wilding, H.M.V., Takahashi, E.A., Newell, M., Wanless, S., & Cunningham, E.J.A. (2012). Impacts of parasites in early life: contrasting effects on juvenile growth for different family members. *PLoS One*, 7, e32236.
- RENTAS. (2014). 1º Relatório Nacional sobre o tráfico de fauna silvestre. Brasil. [https://renctas.org.br/wp-content/uploads/2014/02/REL\\_RENCTAS\\_pt\\_final.pdf](https://renctas.org.br/wp-content/uploads/2014/02/REL_RENCTAS_pt_final.pdf)
- Ribeiro, L.B., & Silva, M.G. (2007). O comércio ilegal põe em risco a diversidade das aves no Brasil. *Ciência e Cultura*, 59, 4-5.
- Ryan, U. (2010). *Cryptosporidium* in birds, fish and amphibians. *Experimental parasitology*, 124, 113-120.
- Saegerman, C., Dal Pozzo, F., & Humblet, M.F. (2012). Reducing hazards for humans from animals: emerging and re-emerging zoonoses. *Italian Journal of Public Health*, 9, 13-24.
- Santos, I.G., Batista, A.I.V., Da Silva, W.S.I., Oliveira Neto, M.B., Schettino, S.C., Oliveira, M.R., Ramos R.A.N., Alves, L.C., Bezerra-Santos, M., & Lima, V.F.S. (2022). Gastrointestinal parasites in captive wild animals from two Brazilian Zoological Gardens. *Research, Society and Development*, 11, e28411426637.
- Silva, G., Alves, L., Borges, J.C.G., Faustino, M.A.G. (2016). Ocorrência de *Cryptosporidium* spp. em felinos de Recife, PE, Brasil. *Veterinária e Zootecnia*, 22, 408-417.
- Snak, A., Lenzi, P.F., Agostini, K.M., Delgado, L.E., Montanucci, C.R., & Zabott, M.V. (2014). Análises coproparasitológicas de aves silvestres cativas. *Ciência Animal Brasileira*, 15, 502-507.
- Sousa, T.N., Sousa, R.V.S., Evangelista, B.B.C., & Freire, S.M. (2018). Prevalência das zoonoses parasitárias e a sua relação com as aves silvestres no Nordeste do Brasil. *Jornal Interdisciplinar de Biociências*, 3, 39-44.
- Souza, L.S., Andrade, A.M.F., Guilherme, E., & Santos, F.G.A. (2019). Endoparasites in wild birds in the Brazilian Amazon. *Brazilian Journal of Veterinary Medicine*, 41, e105219.
- Sprenger, L.K., Yoshitani, U.Y., Buzatti, A., Molento, M.B. (2018). Occurrence of gastrointestinal parasites in wild animals in State of Paraná, Brazil. *Anais da Academia Brasileira de Ciências*, 90, 231-238.

Received June 2, 2024.

Accepted August 2, 2024.