




Neotropical Helminthology



ORIGINAL ARTICLE / ARTÍCULO ORIGINAL

BIODIVERSITY OF PARASITES INFECTING *BOTHROPS ERYTHROMELAS* AMARAL, 1923 (SQUAMATA, VIPERIDAE): AN ENDEMIC VENOMOUS SNAKE SPECIES FROM THE BRAZILIAN NORTHEAST

BIODIVERSIDAD DE PARÁSITOS QUE INFECTAN A *BOTHROPS ERYTHROMELAS* AMARAL, 1923 (SQUAMATA, VIPERIDAE): UNA ESPECIE DE SERPIENTE VENENOSA ENDÉMICA DEL NORESTE DE BRASIL

Cicera Silvilene Leite Matias^{1,2,*}; Cristiana Ferreira-Silva³; Willianilson Pessoa-da-Silva^{1,2}; Maria Beatriz de Andrade Sousa^{1,2}; Aldenir Ferreira Silva-Neta²; Robson Waldemar Ávila³; Felipe Gobbi Grazziotin⁴ & Adrian Antonio Garda^{1,2}

- ¹ Programa de Pós-graduação em Ciências Biológicas (Zoologia) Universidade Federal da Paraíba - Via Expressa Padre Zé Jardim, Cidade Universitária, 58051001, João Pessoa, PB, Brazil;
- ² Laboratório de Anfíbios e Répteis, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Campus Universitário, Lagoa Nova, Av. Senador Salgado Filho, 3000, 59078-900, Natal, RN - Brasil
- ³ Universidade Federal do Ceará (UFC), Departamento de Biologia, Centro de Ciências, Av. Mister Hull, s/n, CEP 60455-760, Fortaleza, Ceará, Brazil
- ⁴ Laboratório de Coleções Zoológicas, Instituto Butantan, São Paulo, 05503-900, SP, Brazil
- * Corresponding author: silvilenematias@gmail.com

Cicera Silvilene Leite Matias: <https://orcid.org/0000-0002-2812-0106>

Cristiana Ferreira-Silva: <https://orcid.org/0000-0002-3087-2614>

Willianilson Pessoa-da-Silva: <https://orcid.org/0000-0002-4975-5733>

Maria Beatriz de Andrade Sousa: <https://orcid.org/0009-0007-9158-8928>

Aldenir Ferreira Silva-Neta: <https://orcid.org/0000-0003-4245-5233>

Robson Waldemar Ávila: <https://orcid.org/0000-0003-3641-8321>

Felipe Gobbi Grazziotin: <https://orcid.org/0000-0001-9896-9722>

Adrian Antonio Garda: <https://orcid.org/0000-0002-1178-1207>

ABSTRACT

The knowledge of parasitic fauna in wild animals is essential for understanding the ecological conditions that determine the occurrence and prevalence of parasites in their hosts. Except for records of one Pentastomida (*Cephalobaena tetrapoda* Heymons, 1922) and two Nematoda (*Physaloptera* sp. And *Aspicularis* sp.), detailed information about the helminth fauna associated with the jararaca *Bothrops erythromelas* Amaral, 1923 is lacking. This species has a wide distribution in the Caatinga, with records in marginal areas of the Cerrado and Atlantic Forest. Here, we describe the patterns of

Este artículo es publicado por la revista Neotropical Helminthology de la Facultad de Ciencias Naturales y Matemática, Universidad Nacional Federico Villarreal, Lima, Perú auspiciado por la Asociación Peruana de Helminthología e Invertebrados Afines (APHIA). Este es un artículo de acceso abierto, distribuido bajo los términos de la licencia Creative Commons Atribución 4.0 Internacional (CC BY 4.0) [<https://creativecommons.org/licenses/by/4.0/deed.es>] que permite el uso, distribución y reproducción en cualquier medio, siempre que la obra original sea debidamente citada de su fuente original.



DOI: <https://dx.doi.org/10.62429/rnh20242181808>

richness, abundance, and prevalence of helminths in *B. erythromelas*, a venomous snake from northeastern Brazil. The parasitized snakes were collected from six Brazilian states in the Northeast region: Bahia, Ceará, Paraíba, Pernambuco, Piauí, and Rio Grande do Norte. We examined the gastrointestinal tract of 127 specimens and found 76 individuals infected with at least one endoparasite, represented by 17 *taxa* of helminths: two Acanthocephala, three Cestoda, and 12 Nematoda. With our results, knowledge about hidden biodiversity is expanded, particularly as this is a pioneering study regarding the helminth fauna of *B. erythromelas*. We also describe the occurrence of the species *Physaloptera lutzi* Cristofaro, Guimarães & Rodrigues, 1976, and *Parapharyngodon hispidus* Ferreira *et al.*, 2021 for the first time in snakes.

Keywords. Acanthocephala – Caatinga – Cestoda – Inventory – Nematoda – Parasites

RESUMEN

El conocimiento de la fauna parasitaria en animales silvestres es esencial para comprender las condiciones ecológicas que determinan la ocurrencia y prevalencia de parásitos en sus hospedadores. Con la excepción de registros de un Pentastomida (*Cephalobaena tetrapoda* Heymons, 1922) y dos Nematoda (*Physaloptera* sp. y *Aspicularis* sp.), falta información detallada sobre la fauna de helmintos asociada con la jararaca *Bothrops erythromelas* Amaral, 1923. Esta especie tiene una amplia distribución en la Caatinga, con registros en áreas marginales del Cerrado y la Mata Atlántica. Aquí, describimos los patrones de riqueza, abundancia y prevalencia de helmintos en *B. erythromelas*, una serpiente venenosa del noreste de Brasil. Las serpientes parasitadas fueron recolectadas en seis estados brasileños de la región Nordeste: Bahia, Ceará, Paraíba, Pernambuco, Piauí y Rio Grande do Norte. Examinamos el tracto gastrointestinal de 127 especímenes y encontramos 76 individuos infectados con al menos un endoparásito, representado por 17 taxones de helmintos: dos Acanthocephala, tres Cestoda y 12 Nematoda. Con nuestros resultados, se amplía el conocimiento sobre la biodiversidad oculta, especialmente porque este es un estudio pionero sobre la fauna de helmintos de *B. erythromelas*. También describimos por primera vez la ocurrencia de las especies *Physaloptera lutzi* Cristofaro, Guimarães & Rodrigues, 1976 y *Parapharyngodon hispidus* Ferreira *et al.*, 2021 en serpientes.

Palabras clave: Acanthocephala – Caatinga – Cestoda – Inventario – Nematoda – Parásitos

INTRODUCTION

Parasitological studies in reptiles have expanded in recent years, particularly those involving snakes (Ávila *et al.*, 2013; Kuzmin *et al.*, 2016; Mati *et al.*, 2015; Matias *et al.*, 2018; Quirino *et al.*, 2018; Oliveira *et al.*, 2020, 2023; Araújo *et al.*, 2020; Halán & Kottferová, 2021; Oliveira *et al.*, 2023; Conga *et al.*, 2024), which can serve as definitive or intermediate hosts for various groups of parasites (Araújo *et al.*, 2020; Oliveira *et al.*, 2021; Ferreira-Silva *et al.*, 2022). Understanding the parasitic fauna of wild animals is crucial for comprehending ecological interactions of parasites and their hosts, including natural history, life cycles, and evolutionary aspects of this relationship (Silva, 2008; Matias *et al.*, 2018). Detrimental effects of parasites on host fitness include, among others, the alteration of coloration patterns and, consequently, negative impacts on reproductive success (Schall & Dearing, 1987; Dunlap & Schall, 1995; Levri, 1999).

Given the importance and biological diversity of the genus *Bothrops*, which comprises approximately 63 species worldwide (Uetz *et al.*, 2023), only 14 species of the genus have been studied in the Neotropical Region to understand their helminth fauna: *B. alternatus* Duméril, Bibron & Duméril, 1854, *B. asper* Garman, 1883, *B. atrox* Linnaeus, 1758, *B. barnetti* Parker, 1938, *B. cotiara* Gomes, 1913, *B. erythromelas* Amaral, 1923, *B. insularis* Amaral, 1922, *B. jararaca* Wied-Neuwied, 1824, *B. jararacussu* Lacerda, 1884, *B. lanceolatus* Bonnaterre, 1790, *B. mattogrossensis* Amaral, 1825, *B. moojeni* Hoge, 1966, *B. neuwiedi* Wagler, 1924, *B. pradoi* Hoge, 1948 (Conga *et al.*, 2024), however, these studies are essential for understanding interspecific relationships.

In this study, we present for the first time, a comprehensive analysis of the helminth fauna associated with *Bothrops erythromelas*, the Caatinga lancehead (jararaca-da-seca in Portuguese). *Bothrops erythromelas* is widely distributed in the Caatinga, throughout the Brazilian states of

Alagoas, Bahia, Ceará, Maranhão, Minas Gerais, Paraíba, Pernambuco, Piauí, Rio Grande do Norte, and Sergipe; with records in marginal areas along the Cerrado and Atlantic Forest ecotones (Campbell & Lamar, 2004; Nogueira *et al.*, 2019; Costa *et al.*, 2023). *Bothrops erythromelas* is the most important species from an epidemiological perspective, as it causes most of the life-threatening snake bites in the Brazilian semi-arid region (Lira-da-Silva *et al.*, 2009). Additionally, the species has a key ecological role, contributing to the population size equilibrium of several small vertebrates in the ecosystem (e.g., rodents, amphibians, lizards; Martins *et al.*, 2002). Currently, only three records of endoparasites were registered for *B. erythromelas*, one Pentastomida (*Cephalobaena tetrapoda* Heymons, 1922; Oliveira *et al.*, 2015) and two Nematoda (*Physaloptera* sp., Oliveira *et al.*, 2018; *Aspicularis* sp., Batista *et al.*, 2021). Therefore, a proper appreciation of the species interaction with parasites is crucial for understanding the occurrence and abundance patterns of endoparasites associated with this venomous snake, as well as the resulting medical and ecological implications (Jorge *et al.*, 2015).

MATERIALS AND METHODS

These parasitized snakes were sourced from four northeastern Brazilian states: Bahia (N = 7), Ceará (N = 33), Paraíba (N = 3), Pernambuco (N = 11), Piauí (N = 1), and Rio Grande do Norte (N = 21). Specimens utilized in this investigation were retrieved from Biological Collections: specifically, the Herpetological Collection of the Regional University of Cariri - URCA (N = 20), the Herpetological Collection of the Semiarid - UFERSA (N = 8), the Herpetological Collection of the Federal University of Ceará - NUROF/UFC (N = 23), and the Herpetological Collection of the Federal University of Paraíba - UFPB (N = 25).

We carefully examined the presence of parasites in the gastrointestinal tract (esophagus, stomach, small and large intestines), liver, kidneys, lungs, gallbladder, and celomic cavity of the snakes under a stereomicroscope. Helminths were identified to the lowest possible taxonomic level. To achieve this, we stained Cestodes and Acanthocephala with alcoholic chloride carmine solution, and cleared with eugenol, while species of Nematoda were cleared in lactic acid (Amato & Amato, 2010). To visualize the internal and external morphological characters of the helminths, we utilized an optical microscope. Following identification, all helminths were deposited in the


Parasitological Collection of the Federal University of Ceara (UFC-P).

We calculated the parasitological parameters of abundance, richness, and prevalence following the method outlined by Bush *et al.* (1997), using the software Quantitative Parasitology 3.0. Richness was determined as the total number of parasite species, while abundance represented the total count of parasite individuals found in each of the 76 specimens of *B. erythromelas*. To assess whether parasite richness is influenced by the linear measure of host size (snout-vent length: SVL), sex, and whether abundance is influenced by SVL, sex, and host maturity, we employed a Generalized Linear Model (GLM). In this analysis, predictor variables were included with interaction terms. Given that richness and abundance data are count data, we incorporated a Poisson error structure into the model. These statistical analyses were conducted using R version 4.3.1 (R Core Team, 2023).

Ethic aspects: The specimens used in this study were collected under the approved authorization (29613-1) conceived by Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio/SISBIO) from the Ministério do Meio Ambiente (MMA) of Brazil. The collecting, hold and storage were previously revised by the Ethics and Animal Welfare Committee from the Universidade Regional do Cariri (#00026/2015).


RESULTS

Out of the 127 snakes analyzed, 76 specimens were found to harbor parasites, comprising 17 helminth taxa: two Acanthocephala (cystacanths of *Centrorhynchus* sp. and Oligacanthorhynchidae gen. sp.) with a prevalence of 46.90%, three Cestoda (specimens of *Oochoristica* sp. and larvae of Plerocercoid and Cysticercoid) with a prevalence of 2.76%, and 12 Nematoda (*Brevimulticaecum* sp., Cosmocercidae gen. sp., *Hexameta boddaertii* Baird, 1860, *Oswaldocruzia* sp., *Parapharyngodon hispidus* Ferreira *et al.*, 2021, *Physaloptera lutzi* Cristofaro, Guimarães & Rodrigues, 1976, *Physaloptera nordestina* Matias, Morais & Ávila, 2020, *Physaloptera* sp., *Physalopteroides venancioi* Lent *et al.* 1946, *Physocephalus* sp., *Strongyloides ophidiae* Pereira, 1929, and unidentified nematode larvae) with a prevalence of 50.34%. The overall parasite prevalence in *B. erythromelas* was 59.84%, with a mean infection intensity of 13.88 ± 7.0 (range: 1–350) (Table 1, Fig. 1).



(Continued Figure 1)

(Continued Figure 1)



(Continued Figure 1)

(Continued Figure 1)




Figure 1. Helminth species parasitizing *Bothrops erythromelas* - **Acanthocephala:** **a** - *Centrorhynchus* sp., **b** - Oligacanthorhynchidae gen. sp., **Cestoda:** **c** - *Oochoristica* sp. (mature proglottids), **d** - Plerocercoid Larvae, **e** - Cisticercoid Larvae, **Nematoda:** **f** - *Brevimulticaecum* sp. (anterior end of the larvae), **g** - Cosmoceridae gen. sp. (female adult), **h and i** - *Hexametra boddaertii* (anterior end and tail of the male, respectively), **j** and **k** - *Oswaldocruzia* sp. (anterior end and posterior end of the female, respectively), **l** and **m** - *Parapharyngodon hispidus* (anterior end and tail of the male, respectively), **n** and **o** - *Physaloptera lutzi* (anterior end and tail of the male, respectively) and **p** - *Physaloptera nordestina* (male's tail), **q** - *Physaloptera* sp., **r** - *Physalopteroides venancioi* (anterior end of the female), **s** and **t** - *Physocephalus* sp. (anterior end and posterior end of the larvae, respectively), **u** and **v** - *Strongyloides ophidiae* (panoramic view and eggs of the female, respectively), **w** and **x** - Nematoda not identified (anterior end and posterior end of the larvae, respectively).

Table 1. Parasitological indices of endoparasites in *Bothrops erythromelas* (equivalent to hosts infected with at least one species of parasite). Development stage (DS), Larvae (L), Adut (A), Prevalence values (P%), number of endoparasites (NE), mean intensity of infection and standard deviation (MII \pm SD), mean abundance (MA), site of infection (SI) Sto= stomach; Igi= large intestine; Smi= small intestine, Per= Peritoneum. Northeastern Brazilian state where the host was collected: Bahia (BA), Ceará (CE), Paraíba (PB), Pernambuco (PE), Piauí (PI), and Rio Grande do Norte (RN). * New record for *B. erythromelas* and ** new record for snakes.

Endoparasites	DS	P%	NE	MII \pm SD	MA	SI	States
Acanthocephala							
<i>Centrorhynchus</i> sp.*	L	13.2	56	5.6 \pm 2.5	25	Per, Sto	CE, PE, RN
Oligacanthorhynchidae gen. sp.*	L	75.0	350	6.1 \pm 4.0	46	Per, Sto, Igi, Smi	BA, CE, PB, PE, PI, RN
Cestoda							
<i>Oochoristica</i> sp.*	A	2.6	3	1.5 \pm 1.5	1	Igi, Smi	BA, CE
Plerocercoid larvae*	L	1.3	2	2.0 \pm 2.0	-	Per	RN
Cisticercoid larvae*	L	1.3	13	13.0 \pm 13.0	-	Sto	CE
Nematoda							
<i>Brevimulticaecum</i> sp.*	L	1.3	1	1.0 \pm 1.0	-	Per	CE
Cosmocercidae gen. sp.*	A	2.6	9	4.5 \pm 4.5	7	Smi, Per	CE, RN
<i>Hexametra boddaertii</i> *	A	9.2	22	3.1 \pm 1.0	9	Per, Sto, Igi	CE, RN
<i>Oswaldocruzia</i> sp.*	A	1.3	1	1.0 \pm 1.0	-	Sto	CE
<i>Parapharyngodon hispidus</i> **	A	1.3	8	8.0 \pm 8.0	-	Sto	CE
<i>Physaloptera lutzi</i> **	A	13.2	41	4.1 \pm 3.5	13	Per, Igi, Sto, Smi	CE, RN
<i>Physaloptera nordestina</i> *	A	2.6	3	1.5 \pm 1.5	1	Per, Sto	PE, RN
<i>Physaloptera</i> sp.	L	22.4	239	14.1 \pm 9.0	65	Smi, Sto, Per, Igi	BA, CE, PE, RN
<i>Physalopteroides venancioi</i> *	A	2.6	16	8.0 \pm 8.0	6	Igi, Sto	BA, CE
<i>Physocephalus</i> sp.*	L	1.3	1	1.0 \pm 1.0	-	Sto	PE
<i>Strongyloides ophidiaie</i> *	A	7.9	217	36.2 \pm 30.5	80	Sto, Per, Smi,	CE, RN
Nematoda (not identified)	L	22.4	73	4.3 \pm 2.0	17	Per, Smi, Igi	BA, CE, PE, RN

Among the parasites found, the cystacanth of Oligacanthorhynchidae was the most abundant, with 350 individuals, followed by *Physaloptera* sp., with 239. The rarest species, with only one individual found, were *Brevimulticaecum* sp., *Oswaldocruzia* sp., and *Physocephalus* sp. Parasite richness was positively

influenced by the SVL of the hosts, with no effect of sex or interaction between them (Table 2, Fig. 2). Parasite abundance was also positively influenced by SVL. Additionally, females tended to have higher parasite abundance than males, and both sexes tended to have higher abundance with increasing SVL (Table 3, Fig. 3).

Table 2. GLM model testing how parasite richness varies as a function of host body condition index (SVL), sex, and their interactions.

Variables	Estimate	Standard error	value Z	Value P
Intercepto	-0.468	0.555	-0.844	0.398
SVL	0.003	0.001	2.188	0.028
Sex	0.364	1.009	0.361	0.718
SVL:Sex	-0.001	0.002	-0.446	0.655




Figure 2. Species richness of parasites in relation to the body condition index (SVL) of *Bothrops erythromelas* specimens.

Table 3. GLM model testing how parasite abundance varies as a function of host body condition index (SVL), sex, and their interactions.

Variables	Estimate	Standard error	Value Z	Value P
Intercept	0.564	0.202	2.788	0.005
SVL	0.005	0.000	11.849	<0.001
Sex	-0.034	0.438	-0.078	0.938
SVL: Sex	0.000	0.001	-0.255	0.798




Figure 3. Parasite abundances in relation to the body condition index (SVL) for males and females of *Bothrops erythromelas* species.

DISCUSSION

Bothrops erythromelas exhibited a high prevalence of helminths (59.84%), similar to other species of snakes in the Viperidae family: *Bothrops moojeni* N = 50 (68%; Barrella & Silva, 2003) and *Crotalus durissus terrificus* Linnaeus, 1758 N = 12 (66%; Dias *et al.*, 2004), but contrasting with *Crotalus tzabcan* Klauber, 1952 N = 50 (14%; Carbajal-Márquez *et al.*, 2018). These comparative data support the idea that describing the helminths of a particular species is crucial to achieve a more complete comprehension of hidden patterns of parasite biodiversity.

The phylum Acanthocephala exhibits an indirect life cycle, engaging in trophic interactions with both invertebrates and vertebrates (Crompton & Nickol, 1985). Incidentally, snakes act as paratenic hosts for acanthocephalans (Travassos, 1917; Pizzatto & Madi, 2002; Pizzatto & Marques, 2006; Smales, 2007a; Matias *et al.*, 2018). In these cases, encysted larvae reside in the body cavity of snakes until they are ingested by the definitive host (Nickol, 1985). Typically, birds and mammals serve as definitive hosts, arthropods act as intermediate hosts, and snakes serve as paratenic hosts of the family Oligacanthorhynchidae (Schmidt, 1972; Nickol & Dunagan, 1989; Kennedy, 2006). Birds and mammals are potential predators of snakes, (Cardoso & Santos, 2012; Fraga *et al.*, 2013; Marques & Medeiros, 2018), including those of the genus *Bothrops* (Sazima, 1922; Oliveira & Santori, 1999; Martins *et al.*, 2003). Thus, we suggest that parasites may use snakes to reach their definitive hosts. Representatives of Oligacanthorhynchidae are known to infect snakes such as *Boa constrictor* Linnaeus, 1758, *Boiga dendrophila* Boie, 1827, *Bothrops jararacussu*, *Bothrops neuwiedi*, *Clelia clelia* Daudin, 1803, *Erythrolamprus aesculapii* Linnaeus, 1758, *E. miliaris* Linnaeus, 1758, *E. poecilogyrus* Wied-Neuwied, 1824, *Macrovipera lebetina obtusa* Linnaeus, 1758, *Mastigodryas bifossatus* Raddi, 1820, *Micrurus corallinus* Merrem, 1820, *Philodryas olfersii* Lichtenstein, 1823, *P. patagoniensis* Girard, 1858, *Xenodon merremii* Wagler, 1824, and *X. histricus* Jan, 1863 (Travassos, 1917; Pizzatto & Madi, 2002; Pizzatto & Marques, 2006; Smales, 2007a; Smales *et al.*, 2020).

Besides species of Oligacanthorhynchidae, we also found *Centrorhynchus* sp. in *B. erythromelas*, an acanthocephalan from family Centrorhynchidae. The genus *Centrorhynchus* infects birds of the Falconiformes and Strigiformes orders as definitive hosts and has invertebrates as intermediate hosts, with reptiles and amphibians serving as paratenic hosts (Torres & Puga, 1996). Cystacanths of *Centrorhynchus* have been recorded in snakes of the

genera *Ahaetulla*, *Bothrops*, *Clelia*, *Chironius*, *Dipsa*, *Echiananthera*, *Imantodes*, *Helicops*, *Leptophis*, *Liophis*, *Mastigodryas*, *Tropidonotus*, and *Philodryas* (Travassos, 1926; Vizcaíno, 1993; Smales, 2007a; Smales, 2007b; Lamas & Lunaschi, 2009; Silva & Muller, 2012; Araújo-Filho *et al.*, 2018; Araújo *et al.*, 2020; Yudhana *et al.*, 2023; Conga *et al.*, 2024). In the present study, we only found cystacanths, suggesting that *B. erythromelas* acts as paratenic hosts, and the acanthocephalans use them to reach their definitive hosts, since Falconiformes and Strigiformes have already been reported preying on snakes (Martins *et al.*, 2003; Costa *et al.*, 2009; Cardoso & Santos, 2012). Species-level identification of the acanthocephalans was not possible due to the immature condition of the specimens (cystacanths), which have not yet developed the reproductive system that is fundamental for species identification.

For Cestoda, we identified the presence of *Oochoeristica* sp. and Plerocercoid larvae and cysticercoid. Different species of snakes have been recognized as hosts of Cestoda, such as *Bothrops alternatus*, *B. jararaca*, *B. moojeni*, *Coluber* sp., *Corallus caninus* Linnaeus, 1758, *Erythrolamprus miliaris*, *Micrurus corallinus*, *Philodryas olfersii*, *Pseudoboa nigra* Duméril, Bibron & Duméril, 1854, and *Thamnophis* sp. (Silva *et al.*, 2001; Silva *et al.*, 2006; Chambrier *et al.*, 2010; Matias *et al.*, 2018; Araújo *et al.*, 2020). The genus *Oochoeristica* has been reported in *Echis carinatus* Schneider, 1801, another Viperidae (Farooq *et al.*, 1983). In *B. erythromelas*, Cestoda were found only in early developmental stages, making species-level identification impossible.

We were able to identify most of the nematodes only at the Phylum level (Nematoda, 17 hosts), excepting one identification at the family level (Cosmocercidae gen. sp., 2 hosts), four at the genus level (*Brevimulticaecum* sp., *Oswaldocruzia* sp., *Physocephalus* sp., and *Physaloptera* sp.), due to the immature stage of the parasites, and six at the species level (*Hexametra boddaertii*, *Parapharyngodon hispidus*, *Physaloptera lutzi*, *Physaloptera nordestina*, *Physalopteroides venancioi*, and *Strongyloides ophidiae*). In terrestrial snakes, the richness of Nematoda tends to be higher, as observed in *Crotalus durissus terrificus* (7 spp.; Dias *et al.*, 2004) and *Crotalus mitchelli* Cope, 1861 (5 spp.; Goldberg *et al.*, 2013), and *Pseudoboa nigra* (3 spp.; Matias *et al.*, 2018). Our results corroborate this tendency, since the nematode fauna in *B. erythromelas* includes at least 12 nematode taxa, encompassing distinct species and larval stages. These results support previous studies (Brouat *et al.*, 2007), indicating that the dominance of nematodes in terrestrial snakes may be related to the habitat of their hosts.

The family Cosmocercidae is frequently reported infecting species of amphibians and reptiles (Rizvi, 2009; Ávila & Silva, 2010; Rizvi & Bursey, 2014; Bursey *et al.*, 2015; Araújo *et al.*, 2020; Draghi *et al.*, 2020). Immature individuals of *Brevimulticaecum* sp. have been reported in fish, amphibians, and reptiles (Sprent, 1979; Moravec & Kaiser, 1994; Moravec *et al.*, 1997; Vieira *et al.*, 2010; Silva *et al.*, 2018; Souza *et al.*, 2020). The genus *Oswaldocruzia* is commonly found in amphibians, although it has also been reported in reptiles, with some records for snakes (Durette-Desset *et al.*, 2006; Goldberg & Bursey, 2007; Anderson *et al.*, 2009; Campião *et al.*, 2014; Teles *et al.*, 2015; Oliveira *et al.*, 2017; Oliveira *et al.*, 2019; Flowers & Beane, 2023). *Physocephalus* uses snakes as paratenic hosts, coleopteran insects as intermediate hosts, and mammals as definitive hosts (Ryzhikov, 1952; Kirillov & Kirillova, 2021). The genus *Physocephalus* has been found in the following North American species of Viperidae: *Crotalus atrox* Baird & Girard, 1853, *C. molossus* Baird & Girard, 1853, *C. Pyrrhus* Cope 1866, *C. scutulatus* Kennicott, 1861, and *Sistrurus tergeminus* Say, 1823 (Goldberg *et al.*, 2001; Goldberg *et al.*, 2002; McAllister *et al.*, 2004; Goldberg *et al.*, 2013). Morphologically, it was not possible to identify *Brevimulticaecum* sp. and *Physocephalus* sp. at the species level due to the immature developmental stage of the larvae. Only female specimens were recorded for Cosmocercidae gen. sp. and *Oswaldocruzia* sp., consequently, the identification at species level was not possible because the main morphological characteristics for Cosmocercidae species identification are only present in males.

Nematodes of the genus *Physaloptera* are common in birds, amphibians, mammals, and reptiles (Ortlepp, 1922; Chabaud, 1975; Anderson *et al.*, 2009; Pereira *et al.*, 2012a; Maldonado Jr. *et al.*, 2019). Nine species of *Physaloptera* are known to occur in Brazilian reptiles, with four reported for snakes: *P. monodens* Molin, 1860, *P. liophis* Vicente & Santos, 1974, *P. obtusissima* Molin, 1860, and *P. nordestina* (Matias *et al.*, 2020). Until now, *Physaloptera lutzi* was described only in lizards (Brito *et al.*, 2014; Maia-Carneiro *et al.*, 2018), and this is the first record of the species occurring in a snake. *Bothrops erythromelas* is a generalist in terms of diet, feeding mainly on lizards, frogs, centipedes and mammals (Martins *et al.*, 2002; Silva-Soares *et al.*, 2022). Given that nematodes possess a cuticle resistant to stomach acids, we suggest that these helminths are associated with the ingestion of the host. *Physaloptera nordestina* was already described for the following northeastern Brazilian snakes: *Oxybelis aeneus* Wagler, 1824, *Xenodon merremii*, and *Pseudoboa nigra* (Matias *et al.*, 2020).

Hexametra boddaertii is a common nematode in Squamata (Ávila & Silva, 2010; Bursey & Brooks, 2011), known to infect *Bothrops* sp., *Crotalus durissus*, *Oxyrhopus trigeminus* Duméril, Bibron & Duméril, 1854, *Philodryas baroni* Berg, 1895, *P. patagoniensis*, and *Pseudoboa nigra* (Skrjabin, 1916; Sprent, 1978; Hartdegen & Gamble, 2002; Pinto *et al.*, 2010; Matias *et al.*, 2018). The genus *Parapharyngodon* has a close association with lizards, mainly of the genus *Tropidurus*, with only one species described for amphibians (Ávila & Silva, 2010; Anjos *et al.*, 2013; Araújo-Filho *et al.*, 2015; Ferreira *et al.*, 2021). *Parapharyngodon hispidus* was described in *Tropidurus hispidus* Spix, 1825 (Ferreira *et al.*, 2021), and the present record is the first for snakes. Since only one specimen of *B. erythromelas* was found with *P. hispidus*, we suggest that the infection is accidental, likely due to ingestion during the snake feeding.

Physalopteroides venancioi is the only species of the genus recorded in South America, found mainly in lizards and amphibians (Ávila & Silva, 2010; Campião *et al.*, 2014). We recorded *P. venancioi* in the small intestine of two snakes, suggesting that the infection occurred accidentally, as suggested for the snakes *Platyceps ventromaculatus* Gray 1834, *P. nigra*, and *P. olfersii* (Al-Moussawi, 2016; Matias *et al.*, 2018; Araújo *et al.*, 2020).

The genus *Strongyloides* parasitizes birds, mammals, amphibians, and reptiles (Graham *et al.*, 2024). In response to infection, animals may exhibit diarrhea, bronchopneumonia, alveolar hemorrhage, interstitial inflammation, altered breathing, enterocyte hyperplasia, and ulceration or squamous metaplasia of mucosal epithelium, depending on the worm life stage (Uzal *et al.*, 2016; Thamsborg *et al.*, 2017; Graham *et al.*, 2024). In reptiles, including snakes, *Strongyloides* infections cause respiratory difficulties and pneumonia (Walden *et al.*, 2020; Graham *et al.*, 2024). Five species of *Strongyloides* are documented in snakes: *S. gulae* Little, 1966, *S. serpentis* Little, 1966, *S. ophidiae*, *S. natricis* Navarro & Lluch, 1993, and *S. mirzai* Singh, 1954 (Graham *et al.*, 2024). *Strongyloides ophidiae* has been observed in the following Brazilian snakes: *Crotalus durissus*, *Erythrolamprus miliaris*, *Oxyrhopus guibei* Hoge & Romano, 1977, and *Philodryas olfersii* (Pereira, 1929a; Santos *et al.*, 2009; Mati & Melo, 2014; Mati *et al.*, 2015; Araújo *et al.*, 2020). Transmission of *S. ophidiae* occurs through cutaneous or oral routes, signs of infection are visible after 7 days, and the species has a direct life cycle (Mati & Melo, 2014). This is the first record of *S. ophidiae* acting as a parasite of the species *B. erythromelas*.

The establishment of parasite populations and communities can be influenced by phylogenetic factors or host characteristics (Poulin, 2004; Kamiya *et al.*, 2014a, 2014b). In the present study, the body condition index (SVL) of the snake *B. erythromelas* was positively related to the richness and abundance of helminths. The host's body is considered the habitat of the parasite, such that a larger specimen can provide a greater area for exploration, colonization, and nutrients to endoparasites (Aho, 1990). Another relevant factor is that individuals with greater longevity experience prolonged exposure to parasitic agents (Aho, 1990; Korralo *et al.*, 2007; Pereira *et al.*, 2012b). This same pattern has been documented for other reptiles (Brito *et al.*, 2014; Araújo-Filho *et al.*, 2016; Silva-Neta & Ávila, 2018; Amorim & Ávila, 2019; Teixeira *et al.*, 2018; 2021; Silva *et al.*, 2023).

Of the 30 species of the genus *Bothrops* that occur in Brazil (Costa *et al.*, 2023), 12 (40%) species have been studied regarding their parasites (Conga *et al.*, 2024). The most studied species were: *B. jararaca* (10 studies: Pereira, 1929b; Rodrigues, 1968; Santos & Rolas, 1973; Fabio, 1979; Correa *et al.*, 1990; Vicente *et al.*, 1993; Scherer *et al.*, 1996; Grego *et al.*, 2004; Silva, 2005; Siqueira *et al.*, 2009), *B. moojeni* (7 studies: Travassos, 1926; Pereira, 1929b; Caubisens Poumarau, 1965; Sprent, 1979; Pinto *et al.*, 2012; Müller *et al.*, 2021), and *B. neuwiedi* (6 studies) (Travassos & Freitas, 1941; Ruiz & Leão, 1942; Silva *et al.*, 2001; Barrella & Silva 2003; Pinto *et al.*, 2012; Morais *et al.*, 2017; Müller *et al.*, 2021). This is the third study on the helminth fauna of *B. erythromelas*. Our findings indicate that among all helminth species found in our analysis, only *Physaloptera sp.* is not a new record for *B. erythromelas*, while the species *P. lutzi* and *P. hispidus* are described for the first time in snakes.

Thus, our results expand the knowledge of parasite hidden biodiversity, particularly as this study represents pioneering research on the helminth fauna of *B. erythromelas*, a venomous snake species endemic to the Caatinga.

ACKNOWLEDGEMENTS

We thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for granting a doctoral scholarship, modality GD to CSLM (#141143/2021-5). CFS and RWA thank to CNPq (#150125/2023-2; 307722/2021-0) and thank Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP #FC3-0198-00006.01.00/22) for research funding. FGG

is supported by grants from CNPq (312016/2021-2 and 405518/ 2021-8) and from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP 2022/12660-4). AAG thanks CNPq for financial support through his productivity research grant (307643/2022-0). The research was permitted under license from the Chico Mendes Institute for Biodiversity Conservation (ICMBio) SISBIO (#82300-1).

Author contributions: CRediT (Contributor Roles Taxonomy)

CSLM = Cicera Silvilene Leite Matias

CFS = Cristiana Ferreira-Silva

WPS = Willianilson Pessoa-da-Silva

MBAS = Maria Beatriz Andrade de Sousa

AFSN = Aldenir Ferreira Silva-Neta

RWA = Robson Waldemar Ávila

FGG = Felipe Gobbi Grazziotin

AAG = Adrian Antonio Garda

Conceptualization: CSLM, CFS, FGG, AAG

Data curation: CSLM, CFS, RWA

Formal Analysis: CSLM, WPS, MBAS, AFSN

Funding acquisition: CSLM, FGG, AAG

Investigation: CSLM, CFS, WPS

Methodology: CSLM, CFS

Project administration: CSLM, AAG

Resources: CSLM, AAG

Software: CSLM, MBAS, AFSN

Supervision: AAG

Validation: CSLM, RWA, FGG, AAG

Visualization: CSLM, CFS, RWA, FGG, AAG

Writing – original draft: CSLM

Writing – review & editing: CSLM, CFS, WPS, MBAS, AFSN, RWA, FGG, AAG

BIBLIOGRAPHIC REFERENCES

- Aho, J.M. (1990). Helminth communities of amphibians and reptiles: comparative approaches to understanding patterns and processes. In Esch, G. W.; Busch, A. O. and Aho, J. M. (Eds) *Parasite Communities: Patterns and Processes*. New York, Chapman & Hall, pp.157–195.
- Al-Moussawi, A.A. (2016). The Parasitic Nematode *Physalopteroides venancioi* in the Snake *Platyceps ventromaculatus* (Gray, 1834) in Baghdad City, Central Iraq. *International Journal of Current Microbiology and Applied Sciences*, 5, 350–357.
- Amato, J.F.R., & Amato, S.B. (2010). Técnicas gerais para coleta e preparação de helmintos endoparasitos de aves. In S. Von Matter *et al.* (eds). *Ornitologia e conservação: ciência aplicada, técnicas de pesquisa e levantamento*, 1ª ed. Technical Books, pp. 369–393.
- Amorim, D.M., & Ávila, R.W. (2019). Infection patterns of helminths in *Norops Brasiliensis* (Squamata, Dactyloidae) from a Humid Forest, Northeastern Brazil and their relation with body mass, sex, host size, and season. *Helminthologia*, 56, 168–174.
- Anderson, R.C., Chabaud, A.G., & Willmott, S. (eds) (2009). *Keys to the Nematode Parasites of Vertebrates: archival Volume*. UK.
- Anjos, L.A., Avila, R.W.A., Ribeiro, S.C., Almeida, W.O., & Silva, R.J. (2013). Gastrointestinal nematodes of the lizard *Tropidurus hispidus* (Squamata: Tropiduridae) from a semi-arid region of north-eastern Brazil. *Journal of Helminthology*, 87, 443–449.
- Araújo-Filho, J.A., Brito, S.V., Almeida, W.D.O., Morais, D.H., & Ávila, R.W. (2015). A new species of *Parapharyngodon* (Nematoda: Pharyngodonidae) infecting *Dermatonotus muelleri* (Anura: Microhylidae) from Caatinga, Northeastern Brazil. *Zootaxa*, 4012, 386–390.
- Araújo-Filho, J. A., Brito, S. V., Lima, V. F., Pereira, A. M. A., Mesquita, D. O., Albuquerque, R. L., & Almeida, W. O. (2016). Influence of temporal variation and host condition on helminth abundance in the lizard *Tropidurus hispidus* from north-eastern Brazil. *Journal of Helminthology*, 91, 312–319.
- Araújo-Filho, J.A., Teles, D.A., Teixeira, A.A.M., Sakamoto, A.C.F.L., Pinto, C.L.M., Silva, E.G., & Almeida, W.O. (2018). *Philodryas nattereri* (Run - Snake) Endoparasites. *Herpetological Review*, 49, 526.
- Araújo, K.C., Silva, C.S., Machado, H.T.S., Oliveira, C.R., & Ávila, R.W. (2020). Endoparasites of *Philodryas olfersii* (Lichtenstein, 1823) In restinga environments of the Parnaíba River Delta, Northeastern Brazil. *Neotropical Helminthology*, 14, 129–141.
- Ávila, R.W., Morais, D.H., Anjos, L.A., Almeida, W.O., & Silva, R.J. (2013). Endoparasites infecting the semiaquatic coral snake *Micrurus surinamensis* (Squamata: Elapidae) in the southern amazonian region, Mato Grosso state, Brazil. *Brazilian Journal of Biology*, 73, 645–647.
- Ávila, R.W., & Silva, R.J. (2010). Checklist of helminths from lizards and amphisbaenians (Reptilia, Squamata) of South America. *Journal of Venomous Animals and Toxins including Tropical Diseases*, 16, 543–572.
- Barrella, T.H., & Silva, R.J. (2003). Digenetic trematodes infection in a *Bothrops moojeni* (Viperidae) population from a fauna rescue in Porto Primavera, São Paulo State. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 55, 243–245.
- Batista, A.I.V., Lucena, G.V.C., Nery, T.F.L., Batista, C.C.N., Batista, J.S., Winkeler, I.E., Rolim, C.M. de M., Coelho, W.A.C., Rocha, E.L.B., Lima, V.F.S., & Pereira, J.S. (2021). Gastrointestinal parasites in wild and exotic animals from a Zoobotanical Park in Northeast of Brazil. *Research, Conservation, Education*, 10, 13, e486101321255.
- Brito, S.V., Corso, G., Almeida, A.M., Ferreira, F.S., Almeida, W.O., Anjos, L.A., Mesquita, D.O., & Vasconcellos, A. (2014). Phylogeny and micro-habitats utilized by lizards determine the composition of their endoparasites in the semiarid Caatinga of Northeast Brazil. *Parasitology Research*, 113, 3963–3972.

- Brouat, C., Kane, M., Diouf, M., Bâ, K., Sall-Drame, R., & Duplantier, J.M. (2007). Host ecology and variation in helminth community structure in *Mastomys* rodents from Senegal. *Parasitology*, *134*, 437–450.
- Bush, A.O., Lafferty, K.D., Lotz, J.M., & Shastak, A.W. (1997). Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology*, *83*, 575–583.
- Burse, C.R., Brooks, D.R. (2011): Nematode parasites of Costa Rican snakes (Serpentes) with description of a new species of *Abbreviata* (Physalopteridae). *Comparative Parasitology*, *78*, 333–358.
- Burse, C.R., Goldberg, S.R., & Grismer, L.L. (2015). New specie of *Cosmocercoides* (Nematoda: Cosmocercidae) and other helminths in *Gonocephalus liogaster* (Squamata; Agamidae) from Peninsular Malaysia. *Acta Parasitologica*, *60*, 631–637.
- Campbell, J.A., & Lamar, W.W. (2004). *The Venomous Reptiles of the Western Hemisphere*, Comstock (Cornell University Press).
- Campião, K.M., Morais, D.H., Dias, O.T., Aguiar, A., Toledo, G., Tavares, L.E.R., & Silva, R.J. (2014). Checklist of helminth parasites of amphibians from South America. *Zootaxa*, *3843*, 1–93.
- Carbajal-Márquez, R.A., González-Solís, D., & Cedeño-Vázquez, J.R. (2018). Endoparasites of *Crotalus tzabcan* (Serpentes: Viperidae), with a checklist in rattlesnakes. *Journal of Parasitic Diseases*, *42*, 303–314.
- Cardoso, S.R.T. & Santos, S.M.A. (2012). Observações sobre predação da serpente *Philodryas patagoniensis* por aves. *Biologia Geral e Experimental*, *12*, 7–9.
- Caubisens Poumarau, E.M. (1965). *Catadiscus longicoecalis* nueva especie parasita de ofidios (Trematoda, Paramphistomidae) con una lista de especies del genero *Catadiscus* Cohn, 1904. *Physis*, *25*, 277–282.
- Chabaud, A.G. (1975). *Keys to genera of the order Spirurida*. In Anderson, R.C. et al. (Eds.) *keys to the nematode parasites of vertebrates*: Commonwealth Agricultural Bureaux, Farnham Royal. UK.
- Chambrier, A., Ammann, M., & Scholz, T. (2010). First species of *Ophiotaenia* (Cestoda: Proteocephalidea) from Madagascar: *O. georgievi* n. sp., a parasite of the endemic snake *Leioheterodon geayi* (Colubridae). *Folia Parasitologica*, *57*, 197–205.
- Conga, D., Carlos, S., Oliveira, G., Bezerra, A., Almeida, B., & Pereira, W. (2024). New Records of Helminth Infection in *Bothrops Atrax* Linnaeus, 1758 from Marajó Island-Brazil and a Survey of Parasitic Helminths in *Bothrops* Species (Squamata, Viperidae) from Neotropical Region. *Veterinary Parasitology: Regional Studies and Reports*, *53*, 101060.
- Costa, H.C., Assis, C.L., Werneck, H.A., Nunes, A.V. & Feio, R.N. (2009). Ataque de um falcão acauã (*Herpetotheres cachinnans*) sobre a serpente boipeva (*Xenodon merremii*) na Mata Atlântica de Minas Gerais, sudeste do Brasil. *Revista brasileira de Zociências*, *11*, 171–173.
- Costa, H.C., Guedes, T.B., & Bérnils, R.S. (2023). Lista de répteis do Brasil: padrões e tendências. *Herpetologia Brasileira*, *12*, 1–108.
- Correa, F.M., Paulino, R.C., Buononato, M.A., & Federsoni, P.A. (1990). *Ochetosoma heterocoelium* (Travassos, 1921) (Trematoda: Digenea: Ochetosomatidae) em novo hospedeiro. *Memórias do Instituto Butantan*, *52*, 11–16.
- Crompton, D.W.T., & Nickol, B.B. (1985). *Biology of the Acanthocephala*. Cambridge University Press.
- Dias, R.J.P., Almeida, S.J.B., Prieto, D.B., & Lima, S.S. (2004). Aspectos ecológicos dos nematoides parasitos de *Crotalus durissus terrificus* Laurenti, 1768 (Ophidia, Viperidae), em Juiz de Fora, Minas Gerais, Brasil. *Revista Brasileira de Zociências*, *6*, 231–235.
- Draghi, R., Drago, F.B., & Lunaschi, L.I. (2020). A new species of *Cosmocercoides* (Nematoda; Cosmocercidae) and other helminths in *Leptodactylus latrans* (Anura; Leptodactylidae) from Argentina. *Anais da Academia Brasileira de Ciências*, *92*, 1–14. e20180499

- Dunlap, K.D., & Schall, J.J. (1995). Hormonal alteration and reproductive inhibition in male fence lizards (*Sceloporus occidentalis*) infected with the malarial parasite *Plasmodium mexicanum*. *Physiological Zoology*, 68, 608–621.
- Durette-Desset, M.C., Anjos, L.A., & Vrcibradic, D. (2006). Three new species of the genus *Oswaldocruzia* Travassos, 1917 (Nematoda, Trichostrongylina, Molineoidea) parasites of *Enyalius* spp. (Iguanidae) from Brazil. *Parasite*, 13, 115–125.
- Fabio, S.P. (1979). Considerações sistemáticas sobre algumas espécies do gênero *Styphlodora* Looss, 1899 (Trematoda). *Revista Brasileira de Biologia*, 39, 229–232.
- Farooq, M., Khanum, Z., & Ansar, R. (1983). Helminth parasites of fresh water tortoise and poisonous snake from Sind, Pakistan. *Pakistan Journal of Zoology*, 15, 31–37.
- Ferreira, A.C.S., Vieira, F.M., Da Silva, D.C.S.N., Ribeiro, L.B., Ferreira, J.A., & Muniz-Pereira, L.C. (2021). *Parapharyngodon hispidus* n. sp. (Nematoda: Pharyngodonidae) in *Tropidurus hispidus* (Spix) (Squamata: Tropiduridae) from Caatinga Biome of the Vale do São Francisco, state of Pernambuco, Brazil with a key for the Neotropical species of the genus *Parapharyngodon* Chatterji. *Zootaxa*, 4980, 195200.
- Ferreira-Silva, C., Alcantara, E.P., Ávila, R.W., & Silva, R.J. (2022). First record of *Cosmocerca podicipinus* (Nematoda: Cosmocercidae) parasitizing *Leptodeira annulata* (Serpentes: Dipsadidae) in northeastern Brazil. *Brazilian Journal of Biology*, 82, e265583.
- Flowers, J.R., & Beane, J.C. (2023). Endoparasites of the Corn Snake, *Pantherophis guttatus* (Linnaeus) (Squamata: Colubridae) from North Carolina, U.S.A. *Comparative Parasitology*, 90, 89–94.
- Fraga, R., Lima, A.P., Prudente, A.L.C., & Magnusson, W.E. (2013). *Guia de cobras da região de Manaus - Amazônia Central = Guide to the snakes of the Manaus region - Central Amazonia*. Inpa.
- Goldberg, S.R., & Bursey, C.R. (2007). *Cercosaura argulus* endoparasites. *Herpetological Review*, 38, 451.
- Goldberg, S.R., Bursey, C.R., & Holycross, A.T. (2001). *Sistrurus catenatus edwardsii* (Desert Massasauga). Endoparasites. *Herpetological Review*, 32, 265.
- Goldberg, S.R., Bursey, C.R., & Painter, C.W. (2002). Helminths of the western diamondback rattlesnake, *Crotalus atrox*, from southeast New Mexico rattlesnake roundups. *Southwestern Naturalist*, 47, 307–310.
- Goldberg, S.R., Bursey, C.R., & Glaudas, X. (2013). Helminths of the Speckled Rattlesnake, *Crotalus mitchellii* (Squamata: Viperidae). *Western North American Naturalist*, 73, 533–535.
- Graham, E.A., Kamp, E.W.L., Thompson, N.M., Tillis, S.B., Childress, A.L., Wellehan, J.F.X., Walden, H.D.S., & Ossiboff, R.J. (2024). Proliferative *strongyloidiasis* in a colony of colubrid snakes. *Veterinary Pathology*, 61, 109–118.
- Grego, K.F., Gardiner, C.H., & Catão-Dias, J.L. (2004). Comparative pathology of parasitic infections in free-ranging and captive pit vipers (*Bothrops jararaca*). *Veterinary Record*, 154, 559–562.
- Halán, M., & Kottferová, L. (2021). Parasitic helminths in snakes from the global legal trade. *Helminthology*, 58, 415–419.
- Hartdegen, R.W., & Gamble, K.C. (2002): *Philodryas baroni baroni* (Baron's racer). Endoparasitism. *Herpetological Review*, 33, 141.
- Jorge, R.J.B., Monteiro, H.S.A., Gonçalves-Machado, L., Guarnieri, M.C., Ximenes, R.M., Borges-Nojosa, D.M., Luna, K.P.O., Zingali, R.B., Corrêa-Netto, C., Gutiérrez, J.M., Sanz, L., Calvete, J.J., & Pla, D. (2015). Venomics and antivenomics of *Bothrops erythromelas* from five geographic populations within the Caatinga ecoregion of northeastern Brazil. *Journal of Proteomics*, 114, 93–111.
- Kamiya, T., O'dwyer, K., Nakagawa, S., & Poulin, R. (2014a). What determines species richness of parasitic organisms. A meta-analysis across animal, plant and fungal hosts. *Biological Reviews*, 89, 123–134.

- Kamiya, T., O'dwyer, K., Nakagawa, S., & Poulin, R. (2014b). Host diversity drives parasite diversity: meta-analytical insights into patterns and causal mechanisms. *Ecography*, *37*, 689–697.
- Korallo, N.P., Vinarski, M.V., Krasnov, B.R., Shenbrot, G.I., Mouillot, D., & Poulin, R. (2007). Are there general rules governing parasite diversity? Small mammalian hosts and gamasid mite assemblages. *Diversity and Distributions*, *13*, 353–360.
- Kennedy, C.R. (2006). Ecology of the Acanthocephala. Cambridge University Press. 1–240.
- Kirillov, A.A., & Kirillova, N.Y. (2021). First finding of spirurid larva (Chromadorea, Spirurida) in the common European viper *Vipera berus* (Linnaeus, 1758) of the Russian fauna. *IOP Conference Series: Earth and Environmental Science*, *818*, 012017.
- Kuzmin, Y., Giese, E.G., Melo, F.T.V., Costa, P.A.F.B., Maschio, G.F., & Santos, G.N.S. (2016). Description of *Serpentirhabdias atroxi* n. sp. (Nematoda: Rhabdiasidae), a parasite of *Bothrops atrox* (Linnaeus) (Reptilia: Serpentes: Viperidae) in Brazilian Amazonia. *Systematic Parasitology*, *93*, 37–45.
- Lamas, M.F., & Lunaschi, L.I. (2009). Primer registro de *Centrorhynchus* sp. (Acanthocephala: Centrorhynchidae) en *Leptophis ahaetulla marginatus* (Colubridae) de Argentina. *Cuadernos de herpetología*, *23*, 45–49.
- Levri, E.P. (1999). Parasite-induced change in host behavior of a freshwater snail: parasitic manipulation or by product of infection? *Behavioral Ecology*, *10*, 234–241.
- Lira-da-Silva, R.M., Mise, Y.F., Casais-e-Silva, L.L., Ulloa, J., Hamdan, B., & Brazil, T.K. (2009). Snakes of medical importance in northeastern of Brazil. *Gazeta Médica da Bahia*, *79*, 7–20.
- Maia-Carneiro, T., Motta-Tavares, T., Ávila, R.W., & Rocha, C.F.D. (2018). Helminth infections in a pair of sympatric congeneric lizard species. *Parasitology Research*, *117*, 89–96.
- Maldonado Jr, A., Simões, R.O., Luiz, J.S., Costa-Neto, S.F., & Vilela, R.V. (2019). A new species of *Physaloptera* (Nematoda: Spirurida) from *Proechimys gardneri* (Rodentia: Echimyidae) from the Amazon rainforest and molecular phylogenetic analyses of the genus. *Journal of Helminthology*, *94*, e68.
- Marques, O.A.V. & Medeiros, C.R. (2018). *Nossas incríveis serpentes: caracterização, biologia, acidentes e conservação*. Cotia: Ponto A.
- Martins, M., Marques, O.A.V., & Sazima, I. (2002). Ecological and phylogenetic correlates of feeding habits in Neotropical pitvipers of the genus *Bothrops*, in Schuett, G.W. et al. (eds) *Biology of the Vipers*, Eagle Mountain Publishing, pp. 307–328.
- Martins, M., Spina, F., Monteiro, C., Sawaya, R.J., & Ariedi-Junior, V.R. (2003): *Bothrops alternatus* (Urutu). Predacion. *Herpetological Review*, *34*, 147–148.
- Mati, V.L.T., & Melo, A. (2014). Some aspects of the life history and morphology of *Strongyloides ophidiae* Pereira, 1929 (Rhabditida: Strongyloididae) in *Liophis miliaris* (Squamata: Dipsadidae). *Neotropical Helminthology*, *8*, 203–216.
- Mati, V.L.T., Pinto, H.A., & Melo, A.L. (2015). Helminths of *Liophis miliaris* (Squamata: Dipsadidae): a list of species and new records. *Helminthology*, *52*, 159–166.
- Matias, C.S.L., Ferreira-Silva, C., Sousa, J.G.G., & Ávila, R.W. (2018). Helminths infecting the black false boa *Pseudoboa nigra* (Squamata Dipsadidae) in northeastern Brazil. *Acta Herpetologica*, *13*, 171–175.
- Matias, C.S.L., Morais, D.H., & Avila, R.W. (2020). *Physaloptera nordestina* n. sp. (Nematoda: Physalopteridae) parasitizing snakes from Northeastern Brazil. *Zootaxa*, *4766*, 173–180.
- McAllister, C.T., Bursey, C.R., & Roberts, J.F. (2004). *Physocephalus sexalatus* (Nematoda: Spirurida: Spirocercidae) in three species of rattlesnakes, *Crotalus atrox*, *Crotalus lepidus* and *Crotalus scutulatus*, from southwestern Texas. *Journal of Herpetological Medicine and Surgery*, *14*, 10–12.

- Morais, D., Aguiar, A., Müller, M., Narciso, R., Da Silva, L., & Da Silva, R. (2017). Morphometric and phylogenetic analyses of *Serpentirhabdias viperidicus* n. sp. (Nematoda: Rhabdiasidae) from the lancehead snake *Bothrops moojeni* Hoge, 1966 (Reptilia: Serpentes: Viperidae) in Brazil. *Journal of Helminthology*, *91*, 360–370.
- Moravec, F., & Kaiser, H. (1994). *Brevimulticaecum* sp. larvae (Nematoda: Anisakidae) from the frog *Hyla minuta* Peters in Trinidad. *Journal of Parasitology*, *80*, 154–156.
- Moravec, F., Prouza, A., & Royero, R. (1997). Some nematodes of freshwater fishes in Venezuela. *Folia Parasitologica*, *44*, 33–47.
- Müller, M.I., Emmerich, E., de Alcantara, E.P., Ungari, L.P., Ebert, M.B., Morais, D.H., O'Dwyer, L.H., & Silva, R.J. (2021). First molecular assessment of two digenean parasites of the lancehead snake *Bothrops moojeni* Hoge, 1966 (Serpentes, Viperidae) in Brazil. *Parasitology Research*, *120*, 971–977.
- Nickol, B.B. (1985). Epizootiology, in Crompton, D.W.T & Nickol, B.B. *Biology of the Acanthocephala*, (eds). Cambridge University Press, pp. 307–346.
- Nickol, B.B., & Dunagan, T. (1989). Reconsideration of the acanthocephalan genus *Echinopardalis*, with a description of adult *E. atrata* and a key to genera of the Oligacanthorhynchidae. *Proceedings of the Helminthological Society of Washington*, *56*, 8–13.
- Nogueira, C.C., Argôlo, A.J.S., Arzamendia, V., Azevedo, J.A., Barbo, F.E., Bérnils, R.S., & Martins, M.C.M. (2019). Atlas of Brazilian snakes: verified point-locality maps to mitigate the Wallacean shortfall in a megadiverse snake fauna. *South American Journal of Herpetology*, *14*, 1–274.
- Oliveira, B.H.S., Teixeira, A.A.N., Queiroz, R.N.M., Araújo-Filho, J.A., Teles, D.A., Brito, S.V., & Mesquita, D.O. (2017). Nematodes infecting *Anotosaura vanzolinia* (Squamata: Gymnophthalmidae) from Caatinga, northeastern Brazil. *Acta Herpetologica*, *12*, 103–108.
- Oliveira, M.C., Lima, V.F., Camilo, I.S., & Almeida, W.O. (2020). *Philodryas olfersii* (Green Snake). Endoparasites. *Herpetological Review*, *51*, 357.
- Oliveira, M.C., Duarte, R.G., Araújo-Filho, J.A., Teles, D.A., & Almeida, W.O. (2021). *Erythrolamprus poecilogyrus* (Green Ground Snake). Endoparasites. *Herpetological Review*, *52*, 665–666.
- Oliveira, M.C., Lima, V.F., Pinto, C.L.M., Silva, É.G., Teles, D.A., Silva, C.F., & Almeida, W.O. (2019). New record of *Physaloptera* sp. (Nematoda: Physalopteridae) parasitizing *Philodryas nattereri* (Ophidia: Dipsadidae) in Brazil. *Herpetology Notes*, *12*, 1031–1034.
- Oliveira, M.C., Lima, V.F., Teles, D.A., Araújo-Filho, J.A., & Almeida, W.O. (2018). *Bothrops erythromelas* (Caatinga Lancehead). Endoparasites. *Herpetological Review*, *49*, 542–543.
- Oliveira, M.C., Martins, A.A., Teles, D.A., Araújo-Filho, J.A., Brito, S.V., & Almeida, W.O. (2015). *Bothrops erythromelas* (Jararaca Dry). Endoparasite. *Herpetological Review*, *46*, 444.
- Oliveira, M.C., Ferreira-Silva, C., Silva, R.J., Almeida, W.O., França, F.G.R., & Lourenço-de-Moraes, R. (2023). First record of *Sebekia oxycephala* (Pentastomida: Sebekidae) infecting *Helicops angulatus* (Reptilia: Colubridae) in Northeast Brazil. *Brazilian Journal of Biology*, *83*, e275302.
- Oliveira, M.E., & Santori, R.T. (1999). Predatory behavior of the opossum *Didelphis albiventris* on the pitviper *Bothrops jararaca*. *Studies on Neotropical Fauna and Environment*, *34*, 72–75.
- Ortlepp, R.J. (1922). The nematode genus *Physaloptera* Rudolphi. *Proceedings of the Zoological Society of London*, *4*, 999–1107.
- Pereira, C. (1929a). *Strongyloides ophidiae*. *Boletim Biológico de São Paulo*, *15*, 16–17.
- Pereira, C. (1929b). Revisão do gênero *Opisthogonimus* (Trematoda). *Revista do Museu Paulista*, *16*, 995–1014.

- Pereira, F.B., Alves, P.V., Rocha, B.M., Souza-Lima, S., & Luque, J.L. (2012a). A new *Physaloptera* (Nematoda: Physalopteridae) parasite of *Tupinambis merianae* (Squamata: Teiidae) from southeastern Brazil. *Journal of Parasitology*, 98, 1227–1235.
- Pereira, F.B., Sousa, B.M., & Lima, S.S. (2012b). Helminth community structure of *Tropidurus torquatus* (Squamata: Tropidurus) in a rocky outcrop area of Minas Gerais state, Southeastern Brazil. *Journal of Parasitology*, 98, 6–10.
- Pinto, H.A., Mati, V.L., & Melo, A.L. (2012). New hosts and localities for trematodes of snakes (Reptilia: Squamata) from Minas Gerais state, Southeastern Brazil. *Comparative Parasitology*, 79, 238–246.
- Pinto, R.M., Muniz-Pereira, L.C., Panizzutti, M.H.M., & Noronha, D. (2010). Nematode parasites of the Neotropical Rattlesnake, *Crotalus durissus* Linnaeus, 1758 (Ophidia, Viperidae) from Brazil: new records and general aspects. *Neotropical Helminthology*, 4, 137–147.
- Pizzatto, L., & Madi, R.R. (2002). *Micrurus corallinus* (Coral Snake). Endoparasites. *Herpetological Review*, 33, 215.
- Pizzatto, L., & Marques, O.A.V. (2006). Interpopulational variation in sexual dimorphism, reproductive output, and parasitism of the water snake *Liophis miliaris* (Colubridae), in the Atlantic Forest of Brazil. *Amphibia-Reptilia*, 27, 37–46.
- Poulin, R. (2004). Macroecological patterns of species richness in parasite assemblages. *Basic and Applied Ecology*, 5, 423–434.
- Quirino, T.F., Ferreira, A.J.M.G., Silva, M.C., Silva, R.J., Morais, D.H., & Ávila, R.W. (2018). New records of Helminths in Reptiles from five states of Brazil. *Brazilian Journal of Biology*, 78(4), 750–754.
- Rizvi, A.N. (2009). Two new species of amphibian nematodes from Bhadra Wildlife Sanctuary, Western Ghats, India. *Zootaxa*, 2013, 58–68.
- Rizvi, A.N., & Bursey, C.R. (2014). *Cosmocercoides himalayan* sp. nov. (Nematoda, Cosmocercidae) in *Duttaphrynus himalayanus* (Amphibia, Anura) from Dehradun (Uttarakhand), India. *Acta Parasitologica*, 59, 80–84.
- R, Development Core Team R. (2023). A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing.
- Rodrigues, H.O. (1968). Novos hospedeiros de *Paradistomum parvissimum* (Travassos, 1918) Travassos, 1919 (Trematoda, Dicrocoelidae). *Atas da Sociedade de Biologia do Rio de Janeiro. Rio de Janeiro*, 11, 167–168.
- Ruiz, J.M., & Leão, A.T. (1942). Notas helmintológicas: Três novas espécies de *Opisthgonimus* parasitas de ofídeos brasileiros (Trematoda: Plagiorchiidae). *Memórias do Instituto Butantan*, 16, 171–185.
- Ryzhikov, K.M. (1952). On the question of reservoir parasitism of *Physocephalus sexalatus* (Molin, 1860) – nematode of swines. *Proceedings of Helminthological Laboratory of Academy of Sciences of the USSR*, 6, 139–141.
- Santos, J.C., & Rolas, F.J. (1973). Sobre alguns cestoides de *Bothrops* e de *Liophis miliaris*. *Atas da Sociedade de Biologia do Rio de Janeiro. Rio de Janeiro*, 17, 35–40.
- Santos, K., Carlos, B., Paduan, K., Kadri, S., Barrella, T., Amarante, M., & Da Silva, R. (2009). Morphological and molecular characterization of *Strongyloides ophidiae* (Nematoda, Strongyloidiidae). *Journal of Helminthology*, 84, 136–142.
- Sazima, I. (1992). Natural History of the Pitviper, *Bothrops jararaca*, in Southeastern Brazil, in: Campbell, J. A., & Brodie, E.D. (Eds.) *Biology of the pitvipers*. 1st Tyler, Texas, EUA: Selva.
- Schall, J.J., & Dearing, M.D. (1987). Malarial parasitism and male competition for mates in the western fence lizard, *Sceloporus occidetalis*. *Oecologia*, 73, 389–392.
- Scherer, P.O., Desiderio, M.H., Serra-Freire, N.M., Cristalli, R., & Freire, L. (1996). *Travassosascaris araujo* (Sprenst, 1978): Um novo hospedeiro. *Parasitology*, 66–67.

- Schmidt, G.D. (1972). Revision of the class Archiacanthocephala Meyer, 1931 (Phylum Acanthocephala), with emphasis on Oliganthorhynchidae Southwell et Macfie, 1925. *Journal of Parasitology*, 58, 290–297.
- Siqueira, L.R., Panizzutti, M.H., Muniz-Pereira, L.C., & Pinto, R.M. (2009). Gross lesions induced by nematodes of *Bothrops jararaca* and *Bothrops alternatus* in Brazil with two new records. *Neotropical Helminthology*, 3, 29–34.
- Silva-Neta, A. F., & Ávila, R. W. (2018). Helminth of the lizard *Colobosauroides cearensis* (Squamata, Gymnophthalmidae) in an area of Caatinga, northeastern Brazil. *Acta Herpetologica*, 13, 95–100.
- Silva, C.S., Ávila, R.W., & Morais, D.H. (2018). Helminth community dynamics in a population of *Pseudopaludicola pocoto* (Leptodactylidae: Leiuperinae) from Northeast-Brazilian. *Helminthology*, 55, 292–305.
- Silva, D.S., & Müller, G. (2012). Primeiro registro de *Opisthogonimus lecithonotus* (Trematoda: Plagiorchiidae) em *Philodryas olfersii* (serpentes: Dipsadidae) e primeiro registro de *P. olfersii* como hospedeiro paratênico de *Centrorhynchus* sp. (Acanthocephala: Centrorhynchidae). *The Biologist (Lima)*, 10, 91.
- Silva, G.D., Teixeira, A.A.M., Franzini, L.D., Mesquita, D.O., & Brito, S.V. (2023). A Temporal variation in diet and helminth abundance in the spiny-tailed lizard, *Strobilurus torquatus* Wiegmann, 1834 (Squamata: Tropiduridae) from the Brazilian Atlantic Forest. *Acta Herpetologica*, 18, 105–114.
- Silva, R.J., (2005). Copulation of *Opisthogonimus fonsceai* Ruiz & Leão, 1942 (Trematoda, Digenea, Plagiorchiidae), parasite of *Bothrops moojeni* Hoge, 1966 (Serpentes, Viperidae). *Journal of Venomous Animals and Toxins including Tropical Diseases*, 11, 68–75.
- Silva, R.J. (2008). Contribuição ao conhecimento da helmintofauna de serpentes da região do Pantanal e entorno, Estado do Mato Grosso do Sul e revisão sobre helmintos parasitas de serpentes no Brasil. [Tese (Livre-docência)]. *Instituto de Biociências de Botucatu, Universidade Estadual Paulista, São Paulo, Brazil*.
- Silva, R.J., Karasawa, A.S., Cherubini, A.L., Barrella, T.H., Lopes, C.A., & Amarante, A.F. (2001). The occurrence of *Crepidobothrium* sp. (Cestoda, Proteocephalidae) in *Bothrops moojeni* (Hoge) (Serpentes, Viperidae). *Brazilian Journal of Biology*, 18, 375–379.
- Silva, R.J., Portela, R.C., & Santos, F.J.M. (2006). *Corallus caninus* (Serpentes, Boidae): a new host for *Ophiotaenia* sp. (Cestoda, Proteocephalidae). *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 98, 961–963.
- Silva-Soares, T., Watanabe, L.K., & Segadilha, J. (2022). Predation of *Scolopendra viridicornis* (Newport 1844) (Chilopoda, Scolopendridae) by *Bothrops erythromelas* (Amaral 1923) (Squamata, Viperidae) in the caatinga, northeast Brazil. *Oecologia Australis*, 26, 60–63.
- Skrjabin, K.I. (1916). Contribution a l'étude de la faune helminthologique du Paraguay. *Zoologicheskii Vestnik*, 1, 705–757.
- Smales, L.R. (2007a). Acanthocephala in amphibians (Anura) and reptiles (Squamata) from Brazil and Paraguay with description of a new species. *Journal of Parasitology*, 93, 392–398.
- Smales, L.R. (2007b). Acanthocephalans of Amphibians and Reptiles (Anura and Squamata) from Ecuador, with the description of *Pandosentis napoensis* n. sp. (Neoechinorhynchidae) from *Hyla fasciata*. *Zootaxa*, 1445, 49–56.
- Smales, L., El-Emarah, G.Y., Essa, I.A.M., Abdulzahra, H.K., & Al-Azizz, S.A. (2020). First Records of *Oligacanthorhynchus* (Oligacanthorhynchidae) from the Honey Badger, *Mellivora capensis wilsoni* (Mustelidae) and the West-Asian Blunt-Nosed Viper *Macrovipera lebetina obtusa* (Viperiidae) from North Basrah, Iraq. *Comparative Parasitology*, 87, 44–48.
- Souza, A.C.F., Gama, C.S., Costa, J.F., Costa, A.L.P., & Viana, D.C. (2020). Índices parasitários de *Brevimulticaecum* sp. (Nematoda: Heterocheilidae) em *Potamotrygon motoro* (Chondrichthyes: Potamotrygonidae) capturados no Arquipélago do Bailique, Macapá. *Ensino de Ciências*, 24, 511–515.
- Sprent, J.F.A. (1978). Ascaridoid nematodes of amphibians and reptiles: *Polydelphis*, *Travassosascaris* n.g. and *Hexametra*. *Journal of Helminthology*, 52, 355–384.

- Sprent, J.F.A. (1979). Ascaridoid nematodes of amphibians and reptiles: *Multicaecum* and *Brevimulticaecum*. *Journal of Helminthology*, 53, 91–116.
- Teixeira, A.A.M., Franzini, L.D., Brito, S.V., Almeida, A.M., & Mesquita, D.O. (2018). Very low prevalence of infection by *Physaloptera lutzii* (Nematoda: Physalopteridae) parasitizing *Kentropyx calcarata* (Squamata: Teiidae), from fragments of Atlantic Forest in Northeast Brazil with a summary of nematodes infecting congeneric species. *Herpetology Notes*, 11, 799–804.
- Teixeira, A.A.M., Sampaio, N.K.S., Araujo-Filho, J.A., Teles, D.A., Almeida, W.O., Mesquita, D.O., & Brito, S.V. (2021). Parasitic infection patterns in *Coleodactylus meridionalis* (Squamata: Sphaerodactylidae) from Atlantic Forest fragments, northeast of the Neotropical Region. *Helminthologia*, 58, 356.
- Teles, D.A., Sousa, J.G.G., Teixeira, A.A.M., Silva, M.C., Oliveira, R.H., Silva, M.R.M., & Ávila, R.W. (2015). Helminths of the frog *Pleurodema diplolister* (Anura, Leiuperidae) from the Caatinga in Pernambuco State, Northeast Brazil. *Brazilian Journal of Biology*, 75, 251–253.
- Thamsborg, S.M., Ketzis, J., & Horii, Y. (2017). *Strongyloides* spp. infections of veterinary importance. *Parasitology*, 144, 274–284.
- Torres, P., & Puga, S. (1996). Occurrence of cystacanths of *Centrorhynchus* sp. (Acanthocephala: Centrorhynchidae) in toads of the genus *Eupsophus* in Chile. *Memórias do Instituto Oswaldo Cruz*, 91, 717–719.
- Travassos, L. (1917). Contribuições para o conhecimento da fauna helmintológica brasileira. VI. Revisão dos acantocefalos brasileiros. Parte I. Fam. Gigantorhynchidae Hamman, 1892. *Memórias do Instituto Oswaldo Cruz*, 9, 5–62.
- Travassos, L. (1926). Contribuições para o conhecimento da fauna helmintológica brasileira. XX. Revisão dos Acanthocephalos brasileiros. Parte II. Familia Echinorhynchidae Hamann, 1892, Sub-fam. Centrorhynchinae Travassos, 1919. *Memórias do Instituto Oswaldo Cruz*, 19, 33–125.
- Travassos, L., & Freitas, J.F. (1941). Relatório da quarta excursão do Instituto Oswaldo Cruz, realizada a zona da Estrada de Ferro Noroeste do Brasil, em agosto e setembro de 1940 Pesquisas parasitológicas. *Memórias do Instituto Oswaldo Cruz*, 35, 705–721.
- Vicente, J.J., Rodrigues, H.O., Gomes, D.C., & Pinto, R.M. (1993). Nematóides do Brasil III. Nematóides de Répteis. *Revista Brasileira de Zoologia*, 10, 19–168.
- Vieira, K.R.I., Vicentina, A.W., Paiva, F., Pozo, C.F., Borges, F.A., Adriano, E.A., Costa, F.E.S., & Tavares, L.E.R. (2010). *Brevimulticaecum* sp. (Nematoda: Heterocheilidae) larvae parasitic in freshwater fish in the Pantanal wetland, Brazil. *Veterinary Parasitology*, 172, 350–354.
- Vizcaíno, S.I. (1993). Presencia del género *Centrorhynchus* Lühe, 1911, (Acanthocephala: Centrorhynchidae) en la República Argentina. *Neotropica*, 39, 77–78.
- Uetz, P., Freed, P., Aguilar, R. & Hošek, J. (2023). The Reptile Database (Accessed July 23, 2023). Electronic database available at <http://www.reptile-database.org/>.
- Uzal, F.A., Plattner, B.L., & Hostetter, J.M. (2016). Alimentary system, in Grant, M, et al. *Pathology of Domestic Animals*. Philadelphia, PA: Saunders, pp. 211–212.
- Walden, H.D., Greiner, E.C., & Jacobson, E.R. (2020). Parasites and parasitic diseases of reptiles, in Jacobson, E.R. *et al. Infectious Diseases and Pathology of Reptiles*, (eds) 2nd. Boca Raton, FL: CRC Press, pp. 859–968.
- Yudhana, A., Praja, R.N., & Edila, R. (2023). First report of acanthocephalan parasite in wild-caught Asian vine snake (*Ahaetulla prasina*) in Indonesia. *Veterinary World*, 16, 317–321.

Received August 9, 2024.

Accepted September 22, 2024.