




# Neotropical Helminthology



## ORIGINAL ARTICLE / ARTÍCULO ORIGINAL

### FIRST REGISTER OF *CONTRACAECUM* SP. AND *EUSTRONGYLIDES* SP. (NEMATODA) IN *KINOSTERNON SCORPIOIDES* (LINNAEUS, 1766) (TESTUDINES: KINOSTERNIDAE) FROM NORTHEASTERN, BRAZIL

### PRIMER REGISTRO DE *CONTRACAECUM* SP. Y *EUSTRONGYLIDES* SP. (NEMATODA) EN *KINOSTERNON SCORPIOIDES* (LINNAEUS, 1766) (TESTUDINES: KINOSTERNIDAE) DEL NORESTE DE BRASIL

Alana Lislea de Sousa<sup>1</sup>, Almerinda Macieira Medeiros<sup>2</sup>, Júlia Boáis Almeida<sup>2</sup>, Silmara Cristina Silva de Aquino<sup>1</sup>, Elias Costa Ferreira Junior<sup>1</sup>, Diego Carvalho Viana<sup>3</sup>, Victor Puicón-Niño de Guzmán<sup>4</sup> & Germán Augusto Murrieta-Morey<sup>1,5\*</sup>

<sup>1</sup> Universidade Estadual do Maranhão (UEMA). Programa de Pós-graduação em Ciência Animal (PPGCA), Cidade Universitária Paulo IV, São Luís, 65055-310, Maranhão-Brasil.

<sup>2</sup> Universidade Estadual do Maranhão (UEMA). Programa de Pós-graduação em Biotecnologia e Biodiversidade da Amazônia Legal (Rede Bionorte), Cidade Universitária Paulo IV, São Luís, 65055-310, Maranhão-Brasil.

<sup>3</sup> Universidade Estadual da Região Tocantina do Maranhão (UEMASUL). Programa de Pós-graduação em Ciência Animal (PPGCA), Cidade Universitária Paulo IV, São Luís, 65055-310, Maranhão-Brasil.

<sup>4</sup> Grupo de Investigación Parasitología Veterinaria y Zoonosis Parasitaria, Laboratorio de Histopatología animal, Facultad de Medicina Veterinaria, Escuela Profesional de Medicina Veterinaria, Universidad Nacional de San Martín. Tarapoto, Perú.

<sup>5</sup> Instituto de Investigaciones de la Amazonia Peruana (IIAP), Laboratorio de Parasitología y Sanidad Acuicola, Carretera Iquitos-Nauta, Km 4.5 – San Juan Bautista, Iquitos, 0784, Loreto, Perú.

\* Corresponding author: germantiss1106@gmail.com

Alana Lislea de Sousa: <https://orcid.org/0000-0002-0920-2560>  
 Almerinda Macieira Medeiros: <https://orcid.org/0000-0002-3808-2035>  
 Júlia Boáis Almeida: <https://orcid.org/0009-0000-0230-732X>  
 Silmara Cristina Silva de Aquino: <https://orcid.org/0000-0001-7680-9699>  
 Elias Costa Ferreira Junior: <https://orcid.org/0000-0002-5925-6372>  
 Diego Carvalho Viana: <https://orcid.org/0000-0002-3302-9892>  
 Victor Puicón Niño de Guzmán: <https://orcid.org/0000-0003-2532-2551>  
 Germán Augusto Murrieta Morey: <https://orcid.org/0000-0001-6244-2654>

## ABSTRACT

*Kinosternon scorpioides* (Linnaeus, 1766) popularly known in Brazil as “Jurará” is a species of chelon that serves as a food resource and source of income for some families in Northeastern Brazil. In that sense, the present study aimed to analyze the metazoan endoparasites infecting *K. scorpioides* from the Baixada Maranhense, Brazil. Fifteen adult specimens

Este artículo es publicado por la revista Neotropical Helminthology de la Facultad de Ciencias Naturales y Matemática, Universidad Nacional Federico Villarreal, Lima, Perú auspiciado por la Asociación Peruana de Helminthología e Invertebrados Afines (APHIA). Este es un artículo de acceso abierto, distribuido bajo los términos de la licencia Creative Commons Atribución 4.0 Internacional (CC BY 4.0) [<https://creativecommons.org/licenses/by/4.0/deed.es>] que permite el uso, distribución y reproducción en cualquier medio, siempre que la obra original sea debidamente citada de su fuente original.



DOI: <https://dx.doi.org/10.62429/rnh20242181853>

were captured using funnel traps in flooded fields around the Aurá River, close to the city of São Bento-Maranhão. Samples were transported to the Postgraduate Multi-User Research Laboratory- Laboratories in Postgraduate Research/ Postgraduate in Animal Science located at the State University of Maranhão- UEMA for parasitological analyses. The results revealed the presence of larvae of *Contraecium* sp. infecting the stomach, intestine, and liver, while *Eustrongylides* sp. was found encysted in the musculature, stomach, intestine, and inside an egg of a gravid female mud turtle. The findings of the present study constitute the first records of these nematodes parasitizing *K. scordioides*. Since *K. scordioides* is consumed as food for some families in the State of Maranhão, Brazil, and the presence of both parasites with hygienic-sanitary importance, complementary studies should be designed to allow us to understand the relationship between host parasites and whether they could have zoonotic relevance.

**Keywords:** Baixada Maranhense – endoparasites – helminths – Jurarã – Nematoda - scorpion mud turtle

## RESUMEN

*Kinosternon scordioides* (Linnaeus, 1766), conocido popularmente en Brasil como “Jurarã”, es una especie de quelonio que sirve como recurso alimenticio y fuente de ingresos para algunas familias del Nordeste de Brasil. En ese sentido, el presente estudio tuvo como objetivo analizar los endoparásitos metazoarios que infectan a *K. scordioides* de la Baixada Maranhense, Brasil. Quince especímenes adultos fueron capturados utilizando trampas de embudo en campos inundados alrededor del río Aurá, cerca de la ciudad de São Bento-Maranhão. Las muestras fueron transportadas al Postgraduate Multi-User Research Laboratory- Laboratories in Postgraduate Research/Postgraduate in Animal Science ubicado en la Universidad Estatal de Maranhão - UEMA para análisis parasitológicos. Los resultados revelaron la presencia de larvas de *Contraecium* sp. infectando el estómago, intestino e hígado, mientras que *Eustrongylides* sp. Se encontró enquistado en la musculatura, estómago, intestino y en el interior de un huevo de una tortuga de lodo hembra grávida. Los hallazgos del presente estudio constituyen los primeros registros de estos nematodos parasitando a *K. scordioides*. Dado que *K. scordioides* es consumido como alimento por algunas familias en el Estado de Maranhão, Brasil, y la presencia de ambos parásitos con importancia higiénico-sanitaria, se deben diseñar estudios complementarios que nos permitan comprender la relación entre los parásitos hospedadores y si pudieran tener relevancia zoonótica.

**Palabras clave:** Baixada Maranhense – endoparásitos – helmintos – Jurarã – Nematoda – tortuga de lodo escorpión

## RESUMO

*Kinosternon scordioides* (Linnaeus, 1766) popularmente conhecido no Brasil como “Jurarã” é uma espécie de quelônio que serve como recurso alimentar e fonte de renda para algumas famílias no Nordeste do Brasil. Nesse sentido, o presente estudo teve como objetivo analisar os endoparasitas metazoários que infectam *K. scordioides* da Baixada Maranhense, Brasil. Quinze espécimes adultos foram capturados usando armadilhas de funil em campos alagados ao redor do Rio Aurá, próximo à cidade de São Bento-Maranhão. As amostras foram transportadas para o Laboratório de Pesquisa Multiusuário de Pós-Graduação - Laboratórios de Pesquisa de Pós-Graduação/Pós-Graduação em Zootecnia localizado na Universidade Estadual do Maranhão - UEMA para análises parasitológicas. Os resultados revelaram a presença de larvas de *Contraecium* sp. infectando o estômago, intestino e fígado, enquanto *Eustrongylides* sp. foi encontrado encistado na musculatura, estômago, intestino e dentro de um ovo de uma tartaruga fêmea grávida. Os achados do presente estudo constituem os primeiros registros desses nematoides parasitando *K. scordioides*. Como *K. scordioides* é consumido como alimento por algumas famílias no estado do Maranhão, Brasil, e a presença de ambos os parasitas com importância higiénico-sanitária, estudos complementares devem ser delineados para nos permitir entender a relação entre os parasitas hospedeiros e se eles podem ter relevância zoonótica.

**Palavras-chave:** Baixada Maranhense – endoparasitas – helmintos – Jurarã – Nematoda – tartaruga-escorpião

## INTRODUCTION

Maranhão is a State that encompasses three of the main Brazilian morphoclimatic domains: Amazon, Caatinga, and Cerrado. The State represents an ecotone zone between these biomes, with 14 distinct vegetation types (IBGE, 2004), reflecting the transition between humid forests and semi-arid habitats (Da Silva, 2011).

The Baixada Maranhense region, placed in Maranhão, is located on the easternmost limit of the Amazon biome, forming part of the Brazilian Legal Amazon. This region is characterized by a landscape made up of lakes, fields, floodplains, igapós, landfills, river-marine systems, and dry land. The floodable natural fields of Baixada Maranhense are extremely complex environments from an ecological point of view, with very diverse structures and functioning (Costa-Neto *et al.*, 2002; Pinheiro, 2013).

Due to the recognized importance of the region's floodable natural fields, the Government of the State of Maranhão established the Baixada Maranhense Environmental Protection Area (APA) with Decree No. 11,900 of June 11, 1991 (State of Maranhão, 1991). In 2000 it was designated by the National Wetlands Committee (CNZU) as a Ramsar Site: regions that provide fundamental ecological services - meeting water and food needs - for species of fauna and flora and human populations, rural and urban. In addition to regulating the water regime of vast regions, they function as a source of biodiversity at all levels. It has the largest set of lake basins in the Northeast, which contribute to high fishing productivity, the main basis of food and income support for its population (Costa-Neto *et al.*, 2002).

Chelonians are animals that perform important ecosystem functions and are intrinsically linked in many traditional cultures, from the consumption and seed dispersal of various tree species to the important social role of communities that use river systems as a means of subsistence (Alves *et al.*, 2008; Rhodin *et al.*, 2018). *Kinosternon scorpioides* (Linnaeus, 1766) popularly known as the scorpion mud turtle and commonly named in Brazil as "Jurará" is a species of chelon widely distributed in the Neotropical region, occurring from Panama, along the Caribbean mountain range to Brazil, covering the entire Amazon forest, eastern mountain ranges of Colombia, Ecuador, and Peru.

Historically, many species of chelonians in different parts of the world have great nutritional, economic, and cultural importance, with their eggs, meat, viscera, fat, and shell being used intensively by humans (Van Dijk *et al.*, 2014). In Baixada Maranhense, species such as *Phrynos geoffroanus* (Schweigger, 1812) and mainly *Kinosternon*

*scorpioides* serve as a food resource and source of income for some families, being captured illegally, in adult and sub-adult form alive and hanging, and sold mainly from May to August, when females appear ovate. Wild populations are used for consumption as a regional delicacy, both for subsistence and recreational consumption (Machado Júnior *et al.*, 2006; Medeiros, 2016).

According to Giari *et al.* (2022) in aquatic environments there exist metazoan parasites with zoonotic potential that can cause strong negative impacts on the safety and quality of the meat. These parasites can be of public health interest and can compromise the health of the consumers. Humans can be infected accidentally with zoonotic parasites by the ingestion of raw or poorly cooked meat (Giari *et al.*, 2022).


Due to the consumption of *K. scorpioides* meat by the Brazilian population and the presence of parasites with zoonotic potential in aquatic organisms, the present study aimed to analyze the metazoan endoparasites infecting *K. scorpioides* from the Baixada Maranhense, Brazil.

## MATERIAL AND METHODS

The study was carried out around the Aurá River, located in the Baixada Maranhense Environmental Protection Area (APA), in the Northeast of Brazil, on the eastern edge of the Amazon (Figure 1). This area represents a wetland of international importance (Ramsar Site), bringing together a set of river basins and flooded fields, and seasonal periods with great contrasts in temperature, humidity, and rainfall (Costa-Neto *et al.*, 2002; Mendes, 2018).

The collections were carried out in flooded fields around the Aurá River, city of São Bento-Maranhão (2°42'08,1"S, 44°51'40,3"W and 2°40'60,0"S, 44°49'00,0"W) (Fig.1). The animals were captured using funnel traps (Secco *et al.*, 2013; Balestra *et al.*, 2016).

Captured animals presented 13,73 cm. mean carapace length, 9,17 cm. mean carapace width, 12,60 cm. mean plastron length, 7,97 cm. mean plastron width, 4,87 cm. mean height and 342,40 g. mean weight. Samples were transported at the Postgraduate Multi-User Research Laboratory- Postgraduate Multi-User Research Laboratory- Laboratories in Postgraduate Research/ Postgraduate in Animal Science located at the State University of Maranhão- UEMA Campus Paulo VI. The samples were collected in the region of the Municipality of São Bento-MA, belonging to the APA of Baixada Maranhense, Sítio Ramsar.



**Figure 1.** Study area. Identification of collection points distributed along the Aurá River (blue) A- Forest areas; B- Flooded fields; C- Water areas (River). Source: IBGE (2018); ZEE (2019); MapBiomias (2021).

The musculature was analyzed and carefully observed. Internal organs were analyzed using a stereoscope Leica EZ4. Organs were placed in Petri dishes with distilled water, and with the aid of tweezers and needles, the tissue was examined by making fine cuts that allowed to observe inside them. In the presence of any parasites, these were removed with tweezers and preserved in ethanol 70% for posterior taxonomic identification.

For taxonomic identification, cysts with nematodes were placed in Petri dishes, and then, with the aid of needles, the cysts were broken, liberating the larvae, and then preserved in 70% ethanol for posterior analyses. For the identification of nematodes, they were placed into glass slides with Lactic Acid, which is used for the clarification of parasite tissue and consequent visualization of external structures and internal organs. All parasites mounted in slides were observed under an optical microscope Leica DM750. Based on the morphological characteristics of the parasites, the identification was made using the information of Moraveck (1998) and Morey *et al.* (2022). Parasitological indices were calculated according to Bush *et al.* (1997).


**Ethic aspects:** The collection of samples was carried out under the permissions granted by the Biodiversity Authorization and Information System (SISBIO), number 85805-1.

## RESULTS


The nematodes *Contracaecum* sp. and *Eustrongylides* sp. were found parasitizing *K. scorpioides*. *Contracaecum* sp. was found in the stomach, intestine, and liver (Fig. 2), while *Eustrongylides* sp. was found encysted in the musculature (Fig. 3) stomach, intestine and inside the egg of a gravid female *K. scorpioides* (Fig. 4). Both parasites were found encysted, being necessary to break the cyst to liberate the larvae. Parasitological indices are presented in Table 1.

*Contracaecum* sp. medium-sized nematodes, opaque-white when alive. Transversely striated cuticles present and more distinct at the extremities of the body, with the anterior region interrupted by a short, lateral line. The cephalic extremity is rounded with a small, ventral cuticular tooth. The oesophageal muscle is narrow, the ventricle is small and rounded, and the ventricular appendix is a short, long intestinal caecum, extending anteriorly to the nerve ring. Tail conical without mucron (Fig. 5A-C).






**Figure 2.** A. Encysted *Contraecum* sp. in internal organs of *Kinosternon scorpioides*. Black circles present the cyst of the parasites. Scale bar: 2 cm.



**Figure 3.** A. *Eustrongylides* sp. in the musculature of *Kinosternon scorpioides*. B. Cyst. C. *Eustrongylides* sp. being released from a cyst. D. Free *Eustrongylides* sp. Scale bar: A, B = 2 cm. C-E = 0.5 cm



**Figure 4.** A. *Eustrongylides* sp. in an egg of *Kinosternon scorpioides*. The black arrow indicates the egg with the parasite. B. Broken egg showing the nematode inside. Scale bar: 2 cm.

**Table 1.** Parasitological indices of endoparasites reported in *Kinosternon scorpioides*. AS = Analyzed specimen. PS = Parasitized specimen. P% = prevalence. TNP = total number of parasites. mI = mean intensity of infection. mA = mean abundance of infection.


Parasite	AS	PS	P%	TNP	mI	mA
<i>Contracaecum</i> sp.	15	11	73.33	19	1.72	0.73
<i>Eustrongylides</i> sp.	15	12	80	32	2.66	2.13

*Eustrongylides* sp. presented a cephalic extremity with a small oral cavity surrounded by 12 cephalic papillae of similar size arranged in two concentric rings, and genital primordia in the posterior part of the body (Fig. 5D-F).


According to the results of the present study, it can be assumed that in the life cycle of *Eustrongylides* sp. and *Contracaecum* sp., different organisms are involved: invertebrates play the role of intermediate hosts, while fish and *K. scorpioides* act as paratenic hosts. Finally, given that *K. scorpioides* are preyed upon by Amazonian alligators, it can be inferred that they act as final hosts for these nematodes in territories of the Baixada Maranhense, Maranhão, Brazil (Fig. 6).

## DISCUSSION

Studies focusing on South American turtles are still recent and scarce (Bursey & Brooks, 2011; Mascarenhas *et al.*, 2013). Few studies have examined the nematodes parasites of *K. scorpioides* in Brazil: *Serpinema magathi* from specimens collected in Pará (Alho, 1965); Freitas and Dobbin Jr. (1971) from Pernambuco; *S. monospiculatus* from specimens collected in a semiarid region in the Northeast (Pereira *et al.*, 2018); *S. pelliculatus* was found from the gastrointestinal tract of specimens collected in Baía do Capim (municipality of Abaetetuba) and Ilha do Marajó (municipality of Soure), Pará, Brazil (Silva *et al.*, 2023). These parasites are commonly found in freshwater turtles worldwide (Hidalgo-Vila *et al.*, 2009; Bursey & Brooks, 2011; Moraga *et al.*, 2012; Mascarenhas *et al.*, 2013; Mascarenhas & Muller, 2015).



**Figure 5.** A. Microphotography of the anterior part of *Contracaecum* sp. Dark circle showing intestinal cecum. B. Microphotography of larval tooth (dark circle). C. Microphotography of the tail of *Contracaecum* sp. D. Encysted *Eustrongylides* sp. E. Microphotography of the anterior part of *Eustrongylides* sp. F. Microphotography of the tail of



**Figure 6.** The life cycle of Nematoda larvae of *Eustrongylides* sp. and *Contracaecum* sp. 1. Nematoda eggs are liberated from the alligator. 2. Eggs with infective L1 larvae. 3. Eggs are consumed by an annelid (for *Eustrongylides* sp.) or for a copepod (for *Contracaecum* sp.). Nematoda larvae develop inside the invertebrates from L1 into L2 and L3. Invertebrates act as intermediate hosts. 4. Fish consume the invertebrates with Nematoda larvae. Fish are paratenic hosts. 5. Fish are consumed by *Kinosternum scorpioides*. The mud turtle acts as a paratenic host. 6. *Kinosternum scorpioides* are consumed by an alligator. The alligator acts as the final host of Nematoda larvae of *Eustrongylides* sp. and *Contracaecum* sp.



For northeastern Brazil, the nematodes *Serpinema magathi* and *Spiroxys figueiredoi* have been found in the digestive tract of *K. scorpioides* (Viana *et al.*, 2016). In the present study, none of these nematodes were found on specimens captured in the municipality of São Bento, Maranhão, Brazil. In the present study, the findings of *Contracaecum* sp. and *Eustrongylides* sp. are the first records of *K. scorpioides*.

Since *K. scorpioides* is an omnivorous species that feeds on algae, plants, seeds, and a variety of insects, mollusks, fish, and small amphibians (Berry & Iverson, 2001; Berry & Iverson, 2011), its feeding habits facilitate the infection of this freshwater turtle by different parasitic nematode larvae present in aquatic environments, such as *Contracaecum* spp. and *Eustrongylides* spp.

*Contracaecum* larvae show low specificity, are often found in abnormal hosts, and do not develop into adults in turtles (Shubber *et al.*, 2020). In this sense, it can be assumed that *K. scorpioides* acquired the nematodes by feeding on fishes in their ecosystems, a fact that was supported by a personal observation made in the field during the capture of the animals in this study.

In the life cycle of *Eustrongylides* spp. freshwater oligochaetes act as intermediate hosts; fish as paratenic hosts and birds as final hosts (Spalding, 2008). *Eustrongylides* spp. can infect poikilothermous hosts, as was demonstrated by experimental infections with piscine, amphibians, and reptilian hosts (Cooper *et al.*, 1978). The presence of nematode larvae in *K. scorpioides* indicates that this turtle acts as the paratenic host of *Contracaecum* sp. and *Eustrongylides* sp. Natural predators of this mud turtle are documented as jaguars (Savage, 2002), owls (Pereira *et al.*, 2007), vultures, green iguanas (Acuña-Mesén, 1998), and alligators (Savage, 2002). The presence of *Eustrongylides* sp. inside the egg of a gravid female *K. scorpioides* can be assumed as a strategy of the parasite to reach easily the final host, since eggs deposited by females are more vulnerable to predators, being predated by birds, terrestrial mammals, and reptiles, facilitating the contact of the nematode with the final host.

In that way, due to the feeding habitat of this chelonian, it is assumed that invertebrates are the intermediate hosts, fish, and *K. scorpioides* paratenic hosts, and alligators are the final hosts.

*Kinosternum scorpioides* from the Baixada Maranhense are parasitized by the nematodes *Contracaecum* sp. and *Eustrongylides* sp. Since *K. scorpioides* is consumed as food for some families in the State of Maranhão, Brazil, and the presence of both parasites with hygienic-sanitary

importance, complementary studies should be designed to allow us to understand the relationship between host parasites and whether they could have zoonotic relevance.

#### Author contributions: CRediT (Contributor Roles Taxonomy)

**ALS** = Alana Lislea de Sousa

**AMM** = Almerinda Macieira Medeiros

**JBA** = Júlia Boáis Almeida

**SCSDA** = Silmara Cristina Silva de Aquino

**ECFJ** = Elias Costa Ferreira Junior

**DCV** = Diego Carvalho Viana

**VPN** = Victor Puicón N.

**GAMM** = Germán Augusto Murrieta Morey

**Conceptualization:** ALS, AMM, JBA, ECF, DCV, ECF, DCV, GAMM

**Data curation:** ALS, GAMM

**Formal Analysis:** ALS, GAMM

**Funding acquisition:** AMM, JBA, SCSDA, GAMM

**Investigation:** SCSDA, ECF, DCV, VPN, GAMM

**Methodology:** ALS, AMM, JBA, SCSDA, ECF, DCV, VPN, GAMM

**Project administration:** ALS, AMM, JBA, GAMM

**Resources:** AMM, JBA, ECF, DCV, GAMM

**Software:** SCSDA, GAMM

**Supervision:** AMM, JBA, GAMM

**Validation:** ALS, AMM, JBA, SCSDA, ECF, DCV, GAMM

**Visualization:** ALS, AMM, JBA, SCSDA, ECF, DCV, VPN, GAMM

**Writing – original draft:** ALS, AMM, JBA, SCSDA, ECF, DCV, VPN, GAMM

**Writing – review & editing:** ALS, AMM, JBA, SCSDA, ECF, DCV, VPN, GAMM

#### ACKNOWLEDGMENTS

We wish to thank the Maranhão State Foundation for Research Development (FAPEMA) for the fellowship granted to SCSDA (002832/2023) and to GAMM (000448/2023).



**BIBLIOGRAPHIC REFERENCES**

- Acuña-Mesen, R.A. (1998). *Las tortugas continentales de Costa Rica*. 2<sup>nd</sup> Ed. 96 pp.
- Alho, C.J.R. (1965). Contribuição ao conhecimento da fauna helmintológica de quelônios do estado do Pará, Brasil. *Boletim do Museu Paraense Emílio Goeldi*, 58, 1–13.
- Alves, R.R.N., Vieira, W.L.S., & Santana, G.G. (2008). Reptiles used in traditional folk medicine: conservation implications. *Biodiversidad y Conservación*, 17, 2037–2049.
- Balestra, R.A.M., Valadao, R.M., & Vogt, R.C. (2016). Roteiro para Inventários e Monitoramentos de Quelônios Continentais. *BioBrasil*, 6, 114–152.
- Berry, J.F., & Iverson, J.B. (2011). *Kinosternon scorpioides* (Linnaeus, 1766) – scorpion mud turtle. In: Rhodin, A.G.F., Pritchard, P.C.H., Van Dijk, P.P., Saumure, R.A., Buhlman, K.A., Iverson, J.B., Mittermeier, R.A. (Eds.), *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtles Specialist Group*. Chelonian Research Monographs, 1–63.
- Berry, J.F., & Iverson, J.B. (2001). *Kinosternon Scorpioides*. Catalogue of American *Amphibians and Reptiles*, 725, 1–11.
- Burse, C.R., & Brooks, D.R. (2011). Nematode parasites of five species of turtles from the Area de Conservación Guanacaste, Costa Rica, with description of a new species of *Falcaustra*. *Comparative Parasitology*, 78, 107–119.
- Bush, A.O., Lafferty, K.D., Lotz, J.M., & Shostak, A. W. (1997). Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology*, 83, 575–583.
- Cooper, C. L., Crites, J. L., & Sprinkle-Fastkie, D. J. (1978). Population biology and behavior of larval *Eustrongylides tubifex* (Nematoda: Diocophymatida) in poikilothermous hosts. *Journal of Parasitology*, 64, 102–107.
- Costa-Neto, J. P., Barbieri, R., Ibañez, M. S. R., Cavalcante, P. R. S., & Piorski, N. M. (2002). Limnologia de três ecossistemas aquáticos característicos da Baixada Maranhense. *Boletim do Laboratório de Hidrobiologia*, 14/15, 19–38.
- Da Silva, A. S. L. (2011). *Aspectos reprodutivos do muçua (Kinosternon scorpioides) em cativeiro* (Tesis de maestría). Universidade Estadual Paulista, Faculdade de Ciências Agrárias e Veterinárias.
- Freitas, J. F. T., & Dobbin Jr., J. E. (1971). Contribuição ao conhecimento da fauna helmintológica de quelônios no estado de Pernambuco, Brasil. *Memórias do Instituto Oswaldo Cruz*, 69, 33–39.
- Giari, L., Castaldelli, G., & Timi, J. T. (2022). Ecology and effects of metazoan parasites of fish in transitional waters. *Parasitology*, 149, 1829–1841.
- Hidalgo-Vila J., Díaz, C. P., Ribas, A. M., Florencio, N., Pérez, S., & Casanova, J. C. (2009). Helminth communities of the exotic introduced turtle *Trachemys scripta* in southwestern Spain: Transmission from native turtles. *Research Journal of Veterinary Science*, 86, 463–465.
- Instituto Brasileiro de Geografia e Estatística (IBGE). (2004). Resolução IBGE R.PR – 1/2004, de 02 de fevereiro de 2019.
- Machado-Júnior, A. A. N., Sousa, A. L., Santos, F. C. F., & Pereira, J. G. (2006). Morfologia dos órgãos genitais femininos do muçua (*Kinosternon scorpioides*). *Arquivo de Ciências Veterinárias*, 11, 25–29.
- Mascarenhas, C. S., & Muller, G. (2015). *Spiroxys contortus* (Gnathostomatidae) and *Falcaustra affinis* (Kathlaniidae) from *Trachemys dorbigni* (Emydidae) in southern Brazil. *Comparative Parasitology*, 82, 129–136.
- Mascarenhas, C. S., Souza, J. D., Coimbra, M. A., & Muller, G. (2013). Nematode parasites of Chelidae (Testudines) from Southern Brazil. *Journal of Parasitology Research*, 112, 3365–3368.
- Medeiros, A. M. (2016). *Aspectos ecológicos, socioambientais e Educação Ambiental aplicados na conservação de Kinosternon scorpioides em comunidades da Baixada Maranhense* (Tesis de maestría). Universidade Estadual do Maranhão.
- Mendes, J. J. (2018). *Dinâmica da paisagem da Bacia do Rio Aurá: um estudo a partir do modelo GTP* (Tesis de maestría). Universidade Estadual do Maranhão.

- Moraga, P., Kinsella, J. M., & Sepulveda, M. S. (2012). Helminth parasites of Eastern box turtles (*Terrapene carolina carolina*) from southern Indiana, USA. *Journal of Helminthology*, 86, 38–40.
- Moravec, F. (1998). *Nematodes of freshwater fishes of the Neotropical Region* (pp. 464). Czech Academy of Sciences.
- Morey, G. A. M., Rojas, C. A. T., Marín, G. A. R., & Guardia, C. T. C. (2022). Occurrence of *Eustrongylides* sp. (Nematoda: Dioctophymatidae) in fish species collected in the Peruvian Amazonia and its implications for public health. *Acta Parasitologica*, 67, 1432–1439.
- Pereira, A. M. A., Brito, S. V., Araújo Filho, J. A., Teixeira, A. A. M., Teles, D. A., Santana, D. O., Lima, V. F., & Almeida, W. O. (2018). Diet and helminth parasites of freshwater turtles *Mesoclemmys tuberculata*, *Phrynops geoffroanus* (Pleurodira: Chelidae) and *Kinosternon scorpioides* (Cryptodira: Kinosternidae) in a semiarid region, Northeast of Brazil. *Acta Herpetologica*, 13, 21–32.
- Pereira, L. A., De Sousa, A.L., Cutrim, M.V.J., & Moreira, E.G. (2007). Características ecológicas do habitat de *Kinosternum scorpioides scorpioides* Linnaeus, 1766 (Reptilia, Chelonia, Kinosternidae) no município de São Bento-Baixada Maranhense (Maranhão, Brasil). *Boletim do Laboratório de Hidrobiologia*, 20, 9-14.
- Pinheiro, C.U.B. (2013). *Matas Ciliares: Recuperação e Conservação em Áreas Úmidas do Maranhão*. São Luis, Maranhão.
- Rhodin, A.G.J., Standford, C.B., & Van Dijk, P.P. (2018). Global Conservation Status of Turtles and Tortoises (Order Testudines). *Chelonian Conservation Biology*, 17, 135–161.
- Salera, J.G., Malvasio, A., & Portelina, T.C.G. (2009). Avaliação da predação de *Podocnemis expansa* e *Podocnemis unifilis* (Testudines, Podocnemididae) no rio Javaés, Tocantins. *Acta Amazonica*, 39, 207–213.
- Savage, J.M. (2002). *The Amphibians and Reptiles of Costa Rica: A Herpetofauna Between Two Continents, Between Two Seas*, Chicago, Illinois. University of Chicago Press, 934 pp.
- State of Maranhao. (1991). Decreto nº 11.900 de 11 de junho de 1991. Dispõe sobre a criação da Área de Proteção Ambiental da Baixada Maranhense no Estado do Maranhão. Publicado no Diário Oficial do Estado nº 195 - em São Luís, 1991.
- Secco, M.C., Mendoca, S.H.S.T., Salles, R.O.L., Silveira, A.L., Luz, V.L.F. (2013). Ocorrência e dados populacionais de *Acanthochelys spixii* na Área de Proteção Ambiental Morro da Pedreira, Serra do Cipó, Minas Gerais, Brasil. In: VI Congresso Brasileiro de Herpetologia. Sociedade Brasileira de Herpetologia.
- Shubber, H. W. K., Alwaaly, A. B. M., & Mohammad, M. K. (2020). Intestinal Helminth Parasites of Caspian Turtle *Mauremys Caspica* (Gmelin, 1774) (Testudines, Geoemydidae) From Al-Diwaniya Province, Iraq. *Annals of Tropical Medicine and Public Health*, 23, 1-11.
- Silva, L. M. O., Jesus, R. F., Santos, A. N., Nandyara, B., Willkens, Y., Santos, J. N., & Melo, F. T. V. (2023). New species of *Serpinema* (Nematoda: Camallanidae) from the scorpion mud turtle *Kinosternon scorpioides* (Testudines: Kinosternidae) from eastern Amazon, Brazil. *International Journal for Parasitology: Parasites and Wildlife*, 22, 6-13.
- Spalding, M., & Forrester, G.J. (2008). *Parasitic diseases of wild birds*. Wiley Blackwell Publishing, Ames.
- Van Dijk, P.P., Iverson, J.B., Rhodin, A.G.J., Shaffer, H.B., & Bour, R. (2014). Turtles of the World. 7th Edition. Annotated checklist of taxonomy, synonymy, distribution with maps and conservation status. Pp. 329–479, In: *Conservation Biology of Fresh Water Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Rhodin, A.G.J., P.C.H. Pritchard, P.P. van Dijk, R.A. Saumure, K.A. Buhlmann, J.B. Iverson, and R.A. Mittermeier (Eds.). Chelonian Research Monographs No. 5.
- Viana, D. C., Rodrigues, J. F. M., Madelaire, C. B., Clara, A., Santos, G., & Sousa, A. L. 2016. Nematoda of *Kinosternon scorpioides* (Testudines: Kinosternidae) from northeastern Brazil. *Journal of Parasitology*, 102, 165-166.
- ZEE. (2019). *Sumário Executivo do Zoneamento Ecológico Econômico do Estado do Maranhão: etapa Bioma Amazônico*. Paulo Henrique de Aragão Catunda; Luiz Jorge Bezerra da Silva Dias (organizadores). São Luís: IMESC.

Received November 5, 2024.

Accepted December 23, 2024.