

ORIGINAL ARTICLE / ARTÍCULO ORIGINAL

EPIDEMIOLOGICAL ASPECTS OF BRONCHOPULMONARY AND GASTROINTESTINAL PARASITE INFECTION IN CATS FROM NORTHEAST BRAZIL

ASPECTOS EPIDEMIOLÓGICOS DE LA INFECCIÓN POR PARÁSITOS BRONCOPULMONARES Y GASTROINTESTINALES EN GATOS DEL NORDESTE DE BRASIL

ASPECTOS EPIDEMIOLÓGICOS DA INFECÇÃO POR PARASITAS BRONCOPULMONARES E GASTROINTESTINAIS EM GATOS DO NORDESTE DO BRASIL

Weslania Souza Inacio da Silva^{1,2}, João Victor Batista dos Santos^{1,2*}, Matheus Resende Oliveira^{1,2,3}, Camenas Vieira Barata^{1,2}, Lorena Maciel Santos Silva^{1,2}, Allan Costa Gomes^{1,2}, Manuel Benicio Oliveira Neto^{1,2}, André Mota Alves^{1,2}, Patrícia Oliveira Meira Santos⁴, Glenda Lídice de Oliveira Cortez Marinho³ & Victor Fernando Santana Lima^{1,2,4}

¹ Program in Parasite Biology, Federal University of Sergipe, São Cristóvão - SE, Brazil.

² Laboratory of Animal Parasitic Diseases, Federal University of Sergipe, Nossa Senhora da Glória - SE, Brazil.

³ Department of Veterinary Medicine, Federal University of Sergipe, Nossa Senhora da Glória.

⁴ Department of Veterinary Medicine, Federal University of Sergipe, São Cristóvão.

* Corresponding author: jvbatista2002@gmail.com

Weslania Souza Inacio da Silva: <https://orcid.org/0000-0003-4393-544X>

João Victor Batista dos Santos: <https://orcid.org/0000-0002-6554-3597>

Matheus Resende de Oliveira: <https://orcid.org/0000-0002-6535-1077>

Camenas Vieira Barata: <https://orcid.org/0009-0002-9312-5963>

Lórena Maciel Santos Silva: <https://orcid.org/0000-0002-3902-278X>

Allan Costa Gomes: <https://orcid.org/0009-0006-5678-8142>

Manuel Benicio Oliveira Neto: <https://orcid.org/0000-0002-7499-4971>

André Mota Alves: <https://orcid.org/0000-0002-9150-5287>

Patrícia Oliveira Meira Santos: <https://orcid.org/0000-0001-6346-3430>

Glenda Lídice de Oliveira Cortez Marinho: <https://orcid.org/0000-0002-7914-9590>

Victor Fernando Santana Lima: <https://orcid.org/0000-0002-7255-0664>

ABSTRACT

This study investigated the presence of bronchopulmonary and gastrointestinal parasites in cats from the state of Sergipe, Brazil, highlighting their medical and veterinary importance. Fecal samples from 100 felines were analysed

Este artículo es publicado por la revista *Neotropical Helminthology* de la Facultad de Ciencias Naturales y Matemática, Universidad Nacional Federico Villarreal, Lima, Perú auspiciado por la Asociación Peruana de Helmintología e Invertebrados Afines (APHIA). Este es un artículo de acceso abierto, distribuido bajo los términos de la licencia Creative Commons Atribución 4.0 Internacional (CC BY 4.0) [<https://creativecommons.org/licenses/by/4.0/deed.es>] que permite el uso, distribución y reproducción en cualquier medio, siempre que la obra original sea debidamente citada de su fuente original.



DOI: <https://doi.org/10.62429/rnh20262012114>

using coproparasitological techniques, showing a 62% positivity rate for parasites, with a significant prevalence of gastrointestinal species. Notably, bronchopulmonary parasites, particularly *Aelurostrongylus* sp., were detected for the first time in Sergipe. Samples were analysed through the Mini-FLOTAC® and Baermann techniques, which proved effective in detecting various parasite types, including eggs, larvae, and cysts. Clinical evaluations were performed, with clinical signs observed in less than 25% of infected cats. The study identified eight parasite genera, with *Ancylostoma* sp. and *Strongyloides* sp. among the most prevalent gastrointestinal parasites, while *Aelurostrongylus* sp. represented bronchopulmonary infections. Both monoparasitism and polyparasitism were recorded, with Mini-FLOTAC® showing superior diagnostic performance. This research presents evidence of *Aelurostrongylus* sp. in domestic cats in Sergipe and underscores the need for enhanced veterinary surveillance. Findings provide a foundation for understanding the transmission dynamics and parasite-host relationships of these diseases, contributing significantly to the epidemiological knowledge in the region.

Keywords: Bronchopulmonary parasites – clinical alterations – feline parasites – Mini-FLOTAC®

RESUMEN

Este estudio investigó la presencia de parásitos broncopulmonares y gastrointestinales en gatos del estado de Sergipe, Brasil, destacando su importancia médica y veterinaria. Se analizaron muestras fecales de 100 felinos mediante técnicas coproparasitológicas, revelando una tasa de positividad del 62% para parásitos, con una prevalencia significativa de especies gastrointestinales. De forma relevante, los parásitos broncopulmonares, particularmente *Aelurostrongylus* sp., fueron detectados por primera vez en Sergipe. Las muestras se analizaron mediante las técnicas Mini-FLOTAC® y Baermann, que demostraron ser eficaces en la detección de diversos tipos de parásitos, incluidos huevos, larvas y quistes. Se realizaron evaluaciones clínicas, observándose signos clínicos en menos del 25% de los gatos infectados. El estudio identificó ocho géneros de parásitos, siendo *Ancylostoma* sp. y *Strongyloides* sp. los parásitos gastrointestinales más prevalentes, mientras que *Aelurostrongylus* sp. representó las infecciones broncopulmonares. Se registraron casos de monoparasitismo y poliparasitismo, presentando la técnica Mini-FLOTAC® un rendimiento diagnóstico superior. Esta investigación aporta evidencias de la presencia de *Aelurostrongylus* sp. en gatos domésticos de Sergipe y refuerza la necesidad de una vigilancia veterinaria más rigurosa. Los resultados proporcionan una base para la comprensión de la dinámica de transmisión y de las relaciones parásito-hospedador de estas enfermedades, contribuyendo de manera significativa al conocimiento epidemiológico de la región.

Palabras clave: Alteraciones clínicas – parásitos broncopulmonares – parásitos de felinos – Mini-FLOTAC®

RESUMO

Este estudo investigou a presença de parasitas broncopulmonares e gastrointestinais em gatos do estado de Sergipe, Brasil, destacando a sua importância médica e veterinária. Amostras fecais de 100 felinos foram analisadas através de técnicas coproparasitológicas, revelando uma taxa de positividade de 62% para parasitas, com prevalência significativa de espécies gastrointestinais. De forma relevante, parasitas broncopulmonares, particularmente *Aelurostrongylus* sp., foram detectados pela primeira vez em Sergipe. As amostras foram analisadas pelas técnicas Mini-FLOTAC® e Baermann, que se mostraram eficazes na detecção de diversos tipos de parasitas, incluindo ovos, larvas e quistos. Foram realizadas avaliações clínicas, tendo sido observados sinais clínicos em menos de 25% dos gatos infetados. O estudo identificou oito gêneros de parasitas, sendo *Ancylostoma* sp. e *Strongyloides* sp. os parasitas gastrointestinais mais prevalentes, enquanto *Aelurostrongylus* sp. representou as infecções broncopulmonares. Registaram-se casos de monoparasitismo e poliparasitismo, com a técnica Mini-FLOTAC® a apresentar desempenho diagnóstico superior. Esta investigação apresenta evidências da presença de *Aelurostrongylus* sp. em gatos domésticos em Sergipe e reforça a necessidade de uma vigilância veterinária mais rigorosa. Os resultados fornecem uma base para a compreensão da dinâmica de transmissão e das relações parasita-hospedeiro destas doenças, contribuindo de forma significativa para o conhecimento epidemiológico na região.

Palavras-chave: Alterações clínicas – parasitos broncopulmonares – parasitos de felinos – Mini-FLOTAC®

INTRODUCTION

In feline medicine, parasitic diseases are increasingly prevalent and considered significant causes of clinical manifestations. Cats can be affected by various gastrointestinal helminths, such as *Ancylostoma* sp., *Toxocara cati* (Schrank, 1788), and *Dipylidium caninum* (Linnaeus, 1758). However, species of nematodes *Ancylostoma* sp. and *Toxocara cati* (Schrank, 1788) are more prevalent (Abbas *et al.*, 2022; Filho *et al.*, 2022; Silva *et al.*, 2023). The diagnosis and study of emerging or neglected parasitic infections have been one of the major challenges in small animal medicine (Abbas *et al.*, 2022).

Bronchopulmonary parasites, for example, have sparked great interest among veterinarians and parasitologists worldwide due to the increase in the number of cases of felines infected with bronchopulmonary nematodes in several European countries and the occurrence of the parasitosis in regions that were previously non-endemic (Traversa & Guglielmini, 2008; Ferraz *et al.*, 2019).

The reasons for the emergence of new cases of bronchopulmonary parasitosis in felines are still unknown; however, some factors such as climate change, intense movement of companion animals, diversity, and susceptibility of potential intermediate hosts (vectors) have been taken into consideration (Livia *et al.*, 2023; Yildirim *et al.*, 2023).

Pulmonary metastrongylids are among the main agents involved in the infection of companion animals, where *Aelurostrongylus abstrusus* (Railliet, 1898) (Nematoda, Metastrongyloidea, Angiostrongylidae), *Troglostrongylus brevior* (Gerichter, 1948), and *Troglostrongylus subcrenatus* (Railliet, 1913) (Nematoda, Strongylida, Crenosomatidae) are listed as the main pulmonary parasites of domestic felines (Livia *et al.*, 2023; Yildirim *et al.*, 2023). The adult forms of these nematodes reside in nodules in the bronchioles and alveolar ducts (Nonnis *et al.*, 2023; Napoli *et al.*, 2023).

As clinical signs in felines infected with bronchopulmonary parasites are not always detectable, as they may present the disease in its asymptomatic or subclinical form. Parasitological examinations, such as the Baermann and Mini-FLOTAC® methods, should be used for confirmation of the diagnosis (Ianniello *et al.*, 2020; Cardoso *et al.*, 2021; Marchiori *et al.*, 2024). However, it's worth noting that young, debilitated, and/or immunosuppressed animals may present with coughing, sneezing, mucopurulent nasal discharge, dyspnea,

tachypnea, exercise intolerance, anorexia, weight loss, and even death (Nonnis *et al.*, 2023). When present, clinical signs result from the intense inflammatory response caused by the elimination of eggs by adult females and the migration of first-stage larvae (L1) to the bronchial tree, causing lesions in the pulmonary alveoli, bronchioles, and local arteries (Vismarra *et al.*, 2023; Yildirim *et al.*, 2023).

Given the impacts that pulmonary and gastrointestinal parasites can have on animal health, the zoonotic potential of some of them, their geographic expansion, and recent advances in epidemiology, clinical, and control of these agents (Abbas *et al.*, 2022), the present study aims to describe the clinical and epidemiological aspects of bronchopulmonary and gastrointestinal parasite infection in naturally infected felines in the state of Sergipe, Brazil.

MATERIALS AND METHODS

Study area

The study included felines residing in rural and/or urban areas of the municipalities of Aracaju, Carira, Nossa Senhora Aparecida, Nossa Senhora das Dores, Nossa Senhora da Glória, and Itabaiana, located in the state of Sergipe, Brazil (10°13'06" S/37°25'13" W) (Figure 1). Sergipe is situated in the Northeast Region and is bordered by the Atlantic Ocean to the east and the states of Bahia to the west and south, and Alagoas to the north, separated by the São Francisco River (IBGE, 2021).

Experimental design and animals

To achieve the proposed objectives, fecal samples were collected from 100 domestic felines of different breeds, sexes, and ages over 2 months old, residing in households, shelters, or Non-Governmental Organizations (NGOs). All animals were clinically evaluated, following three basic steps: 1) Anamnesis and clinical history; 2) Clinical examination; and 3) Fecal sample collection for diagnostic purposes. It is worth noting that all clinical alterations of the animals were described in individual records.

Coproparasitological evaluation

Fecal samples were collected from all felines through spontaneous defecation, with the biological material deposited in properly labeled collection tubes and stored in a cool box at 4°C until further processing at the




Figure 1. Topographic location of the collection areas in the state of Sergipe, Northern Brazil.

Clinical Analysis Laboratory of the Universidade Federal de Sergipe - Sertão campus, Brazil. All samples were analyzed using the Mini-FLOTAC® technique (Cringoli *et al.*, 2013) and the Baermann “gold standard” method (Willesen, 2021).

The Mini-FLOTAC® technique was performed by weighing two grams of feces, which were then deposited into the Fill-FLOTAC® containing 18 mL of zinc sulfate flotation solution ($ZnCl_2$ at 1.350; S7). Subsequently, the process of homogenization and sieving was carried out, and then 5 mL of the solution was transferred to the chamber of the Mini-FLOTAC® until a positive meniscus formed at the end of each chamber. After 10 minutes, the central disk of the Mini-FLOTAC® was rotated at a 90° angle with the aid of a key located at the end of the plate, which was then removed so that the chamber could be attached to the optical microscope adapter. Identification and counting of parasitic structures were then carried out at magnifications of 100X and 400X (Lima *et al.*, 2015).

The Baermann technique was performed by weighing approximately 10g of feces, which were then wrapped in gauze to form a pouch, attached to a rod, and then deposited over a sedimentation glass containing water, ensuring that the fecal sample was fully submerged. After an overnight process, 1.5 mL of sediment from the bottom of the glass was pipetted and deposited into a Falcon tube, which was then centrifuged at a speed of 1500-2000 rpm for 3 min. Following centrifugation, the supernatant was discarded, and approximately 0.1 mL of sediment from the tube was pipetted onto a slide

and cover slip (in triplicate), then examined under an optical microscope at magnifications of 100X and 400X. If necessary, one or two drops of Lugol’s solution were added to fix the larvae (Alho *et al.*, 2013).

All parasites observed in the fecal samples were photographed and analyzed using ImageJ software to obtain morphological and morphometric characteristics, and were then identified based on the taxonomic keys provided by Bowman *et al.* (2006) and Taylor *et al.* (2010).

Statistical analysis

All data were analyzed using InStat software (GraphPad Software), calculating absolute frequency (n/N) and relative frequency (%). The Chi-square test (X^2) was used to determine if there was significance between the diagnostic methods and clinical alterations observed in the animals. The independent variables considered in the model will be those with statistical significance less than 0.20. EpiInfo™7 will also be used to perform statistical calculations, with a significance level established at $p < 0.05$.

Ethical aspects

The present research was submitted to and approved by the Ethics Committee on Animal Use of the Universidade Federal de Sergipe (approval number 4813080621). Animal participation in the research was voluntary, with felines becoming participants after the responsible


guardian signed the Informed Consent Form, following the guidelines outlined in Resolution No. 466/2012 of the National Commission for Ethics in Research of the National Health Council of the Ministry of Health.

RESULTS

In this study, parasitological evaluation revealed positivity for gastrointestinal parasites (93.65%), bronchopulmonary parasites (4.76%), and pseudoparasites (1.59%) in 62.00% (n=62/100; $p < 0.0001$) of fecal samples from

felines in the municipalities of Aracaju (29.03%; 18/62), Nossa Senhora Aparecida (1.61%; 1/62), Nossa Senhora das Dores (3.23%; 2/62), and Nossa Senhora da Glória (66.13%; n=41/62) ($p < 0.0008$).

All biological samples from cats in the municipalities of Carira (0.00%; 0/5) and Itabaiana (0.00%; 0/3) were considered negative. The Venn diagram (Figure 2) schematically demonstrates the presence of bronchopulmonary and gastrointestinal parasites in 16.67% and 66.67% ($p < 0.0001$) of felines from the Sergipe municipalities, respectively.



Legenda: np – Número de parasitos; Ae – *Aelurostrongylus* sp.; An – *Ancylostoma* sp.; Cy – *Cystoisospora* sp.; Dp – *Dipylidium* sp.; Gi – *Giardia* sp.; En – *Entamoeba* sp.; St – *Strongyloides* sp.; To – *Toxocara* sp..

Figure 2. Venn diagram schematically illustrating the presence of bronchopulmonary and gastrointestinal parasites in populations of felines from four municipalities in Sergipe, Northern Brazil.

Parasitic structures were detected in the forms of eggs (59.30%), oocysts (18.60%), cysts (4.65%), larvae (16.28%), and proglottids (1.16%) ($p < 0.0003$) (Figure 2). Eight genera of parasites were identified in the fecal samples of felines (Table 1), with five types of helminths (62.50%) and three protozoa (37.50%) ($p < 0.0026$). Highlighted among the gastrointestinal parasites were *Ancylostoma* sp. (44.00%; 44/62; $p < 0.0011$), *Dipylidium* sp. (1.00%; 1/62; $p < 0.0005$), *Cystoisospora* sp. (19.00%; 19/62; $p < 0.0019$), *Entamoeba* sp. (3.00%;

3/62; $p < 0.0001$), *Giardia* sp. (3.00%; 3/62; $p < 0.0001$), *Strongyloides* sp. (25.00%; 25/62; $p < 0.0063$), and *Toxocara* sp. (2.00%; 2/62; $p < 0.0006$) as the most frequent, followed by the bronchopulmonary parasite *Aelurostrongylus* sp. (3.00%; 3/62; $p < 0.0001$).

Monoparasitism (33.87%; 21/62) was observed exclusively in felines infected only with gastrointestinal parasite genera, while polyparasitism (66.13%; 41/62) ($p < 0.0293$) was identified in animals with two or more

types of enteroparasites, in which symbiotic interactions: bronchopulmonary parasite + gastrointestinal parasite (4.84%; $p < 0.0001$) and gastrointestinal parasite +

gastrointestinal parasite (61.29%; $p < 0.0233$) were reported.

Table 1. Percentage of positivity by type of parasite detected in populations of cats in Northern Brazil.

Parasite/Parasitic structure detected	Absolute Frequency (n/N)	Relative Frequency (%)	Prevalence
<i>Aelurostrongylus</i> sp. ^L	3/62	3.00	3
<i>Ancylostoma</i> sp. ^E	44/62	44.00	44
<i>Cystoisospora</i> sp. ^O	19/62	19.00	19
<i>Dipylidium</i> sp. ^{E,P}	1/62	1.00	1
<i>Entamoeba</i> sp. ^C	3/62	3.00	3
<i>Giardia</i> sp. ^C	3/62	3.00	3
<i>Strongyloides</i> sp ^E	25/62	25.00	25
<i>Toxocara</i> sp. ^E	2/62	2.00	2

Note: E – Eggs, C – Cyst, O – Oocyst, L – Larvae. P – Proglottis

Regarding the effectiveness of the parasitological methods employed, 98.41% and 25.19% of the samples were positive in the Mini-FLOTAC[®] and Baermann techniques, respectively ($p < 0.0001$). 100% of eggs, larvae, cysts, oocysts, and proglottids were diagnosed using the Mini-FLOTAC[®] technique, while the Baermann method was able to recover only *Strongyloides* sp. larvae in 12.50% of the analyzed feline samples ($p < 0.0001$).

Regarding the profile of felines positive for enteroparasites, it was observed that 72.00% were females and 28.00% were males, adults, without defined breed, with an age range between 3 months and 6 years ($p < 0.0153$). Clinically, only 3.23% (2/62) of the animals had a history of respiratory and/or gastroenteritis symptoms in the last 30 days; however, in the clinical examination, 24.19% (15/62) of the felines presented noteworthy alterations ($p < 0.0004$) (Figure 3). Clinical alterations were observed in felines positive for bronchopulmonary and/or gastrointestinal parasites (Figure 4).

DISCUSSION

This study reports for the first time the natural infection of domestic felines by bronchopulmonary parasites in Sergipe, in addition to the description of different gastrointestinal parasites that have an impact on both animal and human health.

The positivity rate (62.00%) for parasitic structures detected in fecal samples of felines from Sergipe was higher than those reported in studies conducted in Australia (8.6%) (Mcglade *et al.*, 2003), China (41.39%) (Yang & Liang, 2015), Egypt (52.40%) (Abbas *et al.*, 2022),

United States (5.1%) (Hoggard *et al.*, 2019), France (Bourgoin *et al.*, 2022), Greece (50.70%) (Symeonidou *et al.*, 2018), Italy (22.00%) (Sauda *et al.*, 2019), India (19.00%) (Vincy & Tresamol, 2023), Russia (18.20%) (Kurnosova *et al.*, 2023), and Zimbabwe (1.1%) (Pfukenyi *et al.*, 2010).

When comparing the overall positivity rate for bronchopulmonary and/or gastrointestinal parasites in patients from our study with published data from epidemiological surveys conducted in Brazil, we observed that studies conducted in Pernambuco using the FLOTAC[®] technique showed a higher percentage of positivity (65.00% to 100%) (Monteiro *et al.*, 2016; Lima *et al.*, 2017). However, research using traditional techniques (e.g., sucrose flotation, spontaneous sedimentation, and direct examination) showed lower positivity rates: Rio de Janeiro (49.50%) (Pereira *et al.*, 2017), São Paulo (18.10%) (Gennari *et al.*, 2016), (26.03%), Goiás (17.00% - 60.00%) (Lima *et al.*, 2018), Rio Grande do Sul (24.63%) (Moreira *et al.*, 2018), and Santa Catarina (37.80%) (Stalliviere *et al.*, 2009).

Clinically, less than 25% of naturally infected felines in Sergipe by bronchopulmonary or gastrointestinal parasites presented clinical alterations. Some authors emphasize that certain parasitized animals may be asymptomatic; however, they may develop cutaneous, gastrointestinal, hepatic, and neurological alterations (Simpson, 1998; Lima *et al.*, 2021). Arslan *et al.* (2019) also highlight that cats are commonly affected by toxocariasis, a gastrointestinal parasitic disease caused by larvae of the ascarid nematodes, *Toxocara canis* (Stiles, 1905) and *T. cati*, which can lead to gastroenteritis, peritonitis,




Figure 3. Parasitic bronchopulmonary and gastrointestinal parasites diagnosed in cats. a - *Ancylostoma* sp. egg; b - *Aelurostrongylus* sp. larva; c - *Cystoisospora* sp. oocyst; d - *Dipylidium* sp. egg capsule; e - *Strongyloides* sp. larvated egg; f - *Toxocara* sp. egg.

eosinophilic ascites due to peritoneal inflammation, abdominal pain, and nausea in infected felines.

In feline medicine, respiratory parasitic infections pose a particular challenge for veterinarians due to the number of pathologies with similar clinical manifestations (Stepanović *et al.*, 2020). Respiratory alterations were observed exclusively in felines parasitized by the bronchopulmonary helminth *Aelurostrongylus* sp., which presented nasal discharge, coughing, sneezing, and

dyspnea, clinical signs that, according to Traversa *et al.* (2010), should be carefully evaluated and considered in the differential diagnosis of feline cardiopulmonary diseases.

Several coproparasitological techniques from the FLOTAC® group are available for the qualitative and/or quantitative diagnosis of gastrointestinal and pulmonary parasitoses in felines. However, some studies highlight the importance of using the Baermann technique in detecting



Figure 4. Clinical alterations observed in felines positive for bronchopulmonary and/or gastrointestinal parasites. a) Animal presenting ocular and nasal discharge; b) Feline presenting cachexia; c) Cat presenting apathy and prostration; and d) Feline presenting weight loss and skin lesions throughout the body.

larvae of bronchopulmonary parasites *Aelurostrongylus abstrusus* and *T. brevior*, which were detected only by the Baermann test (Colombo *et al.*, 2022). Despite the Mini-FLOTAC® technique showing the best performance in diagnosing mixed infections (gastrointestinal and pulmonary), the Baermann test is considered the gold standard for diagnosing cardiopulmonary nematode parasites in animals, due to the positive hydro/thermotropism demonstrated by first-stage larvae present in fecal samples (Morelli *et al.*, 2022).

The findings of this study highlight the presence of bronchopulmonary parasites in the state of Sergipe, as well as the presence of gastrointestinal parasite genera with medical and veterinary significance. Additional studies are warranted to elucidate the parasite-host relationship and the transmission dynamics of these parasitic diseases in Sergipe.

ACKNOWLEDGEMENTS

Thanks to Marcos Antônio Bezerra-Santos the University of Bari and resident of the European Veterinary Parasitology College (EVPC) for contributions to the manuscript. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. The authors are aware of availability of data and materials. The authors declare that the work was carried out without funding.

Author contributions: CRediT (Contributor Roles Taxonomy)

WSIS = Weslania Souza Inacio da Silva

JVBS = João Victor Batista dos Santos

MRO = Matheus Resende Oliveira

CVB = Camenas Vieira Barata

LMSS = Lorena Maciel Santos Silva

ACG = Allan Costa Gomes

MBON = Manuel Benicio Oliveira Neto

AMA = André Mota Alves

POMS = Patrícia Oliveira Meira Santos

GLOCM= Glenda Lídice de Oliveira Cortez Marinho

VFSL=Victor Fernando Santana Lima

Conceptualization: WSIS, VFSL

Data curation: WSIS, JVBS, MRO, CVB, LMSS, ACG, MBON

Formal Analysis: WSIS, JVBS, MRO, MBON

Funding acquisition: WSIS, VFSL

Investigation: WSIS, JVBS, MRO, CVB, LMSS, ACG, MBON

Methodology: WSIS, MRO, MBON

Project administration: POMS, GLOCM, VFSL

Resources: VFSL

Software: WSIS, MRO, MBON, AMA

Supervision: VFSL

Validation: POMS, GLOCM, VFSL

Visualization: WSIS, POMS, GLOCM, VFSL

Writing – original draft: WSIS, JVBS

Writing – review & editing: WSIS, JVBS, AMA, POMS, VFSL

BIBLIOGRAPHIC REFERENCES

- Abbas, I., Al-Araby, M., Elmishmishy, B., & El-Alfy, E. S. (2022). Gastrointestinal parasites of cats in Egypt: high prevalence high zoonotic risk. *BCM Veterinary Research*, 18, 1-12.
- Alho, A. M., Nabais, J., & Carvalho, L. M. (2013). A importância da Técnica de Baermann na clínica de pequenos animais. *Clínica Animal*, 1, 28-31.
- Arslan, F., Baysal, N. B., Aslan, A., Simsek, B. C., & Vahaboglu, H. (2019). *Toxocara* related peritonitis: A case report and review of literature. *International Journal of Parasitology*, 101950, 1-11.
- Bourgoin, G., Callait-Cardinal, M. P., Bouhsira, E., Polack, B., Bourdeau, P., Ariza C.R., Carassou, L., Lienard, E., & Drake, J. (2022). Prevalence of major digestive and respiratory helminths in dogs and cats in France: results of a multicenter study. *Parasites & Vectors*, 15, 1-12.
- Bowman, D. D., Lynn, R. C., Eberhard, M. L., & Alcaraz, A. (2006). *Parasitologia Veterinária de Georgis*. Elsevier.
- Cardoso, B., Pessoa, B., Figueiredo, P., Rinaldi, L., Cringoli, G., Díaz, A., Gomes, L., Santos, N., & Carvalho, L. M. (2021). Comparative survey of gastrointestinal parasites in sympatric Iberian ibex (*Capra pyrenaica*) and domestic goats using molecular host specific identification. *Parasitology Research*, 120, 2291–2296.
- Colombo, M., Morelli, S., Damiani, D., Del-Negro, M. A., Milillo, P., Simonato, G., Barlaam, A., & Di Cesare, A. (2022). Comparison of different copromicroscopic techniques in the diagnosis of intestinal and respiratory parasites of naturally infected dogs and cats. *Animals*, 12, 1-10.
- Cringoli, G., Rinaldi, L., Albonico, M., Berquist, R., & Utzinger, J. (2013). Geospatial (s) tools: integration of advanced epidemiological sampling and novel diagnostics. *Geo Health*, 7, 399-404.
- Ferraz, A., Santos, P. B. S., Santos, E. M., Evaristo, T. A., Coelho, A. L. C., Oliveira, N. L. M., & Nizoli, L. Q. (2019). Verminose pulmonar em felino por *Aelurostrongylus abstrusus* – Relato de caso. *Atas de Saúde Ambiental*, 7, 77-83.
- Filho, C. R. C. U., Santos, K. K. F., Lima, T. A. R. F., Alves, L. C., Carvalho, G. A., & Ramos, R. A. N. (2022). Gastrointestinal parasites in dogs and cats in line with the One Health' approach. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 74, 43-50.
- Gennari, S. M., Ferreira, J. I., Pena, H. F., Labruna, M. B., & Azevedo, S. D. (2016). Frequency of gastrointestinal parasites in cats seen at the University of São Paulo Veterinary Hospital, Brazil. *Revista Brasileira de Parasitologia Veterinária*, 25, 423-428.
- Hoggard, K. R., Jarriel, D. M., Bevelock, T. J., & Verocai, G. G. (2019). Prevalence survey of gastrointestinal and respiratory parasites of shelter cats in northeastern Georgia, USA. *Veterinary Parasitology Regional Studies Reports*, 16, 1-5.

- Ianniello, D., Pepe, P., Alves, L. C., Ciuca, L., Maurelli, M. P., Amadesi, A., Bosco, A., Musella, V., Cringoli, G., & Rinaldi, L. (2020). Why use the Mini-FLOTAC to detect metastrongyloid larvae in dogs and cats. *Acta Parasitologica*, *65*, 546-549.
- IBGE, Instituto Brasileiro de Geografia e estatística (2021). *Sergipe*. <http://www.ibge.gov.br/estadosat/perfil.php?sigla=pe>
- Kurnosova, O. P., Panova, O. A., & Arisov, M. V. (2023). The prevalence of potentially zoonotic intestinal parasites in dogs and cats in Moscow, Russia. *Helminthologia*, *4*, 44-51.
- Lima, J. A. S., Rezende, H. H. A., Rocha, T. M. D. D., & Castro, A. M. (2018). Analysis of the accuracy of different laboratory methods for the diagnosis of intestinal parasites from stray and domiciled cats (*Felis catus domesticus*) in Goiânia, Goiás, Brazil. *Revista Brasileira de Parasitologia Veterinária*, *27*, 95-98.
- Lima, R. L., Pacheco, R. C., Mendonça, A. J., Néspoli, P. E. B., Morita, L. H. M., Almeida, A. D. B. P. F., & Sousa, V. R. F. (2021). *Platynosomum fastosum* in domestic cats in Cuiabá, Midwest region of Brazil. *Veterinary Parasitology Regional Studies Reports*, *24*, 1-4.
- Lima, V. F. S., Ramos, R. A. N., Lepold, R., Borges, J. C. G., Ferreira, C. D., Rinaldi, L., Cringoli, G., & Alves, L. C. (2017). Gastrointestinal parasites in feral cats and rodents from the Fernando de Noronha Archipelago, Brazil. *Revista Brasileira de Parasitologia Veterinária*, *26*, 521-524.
- Lima, V. F. S., Cringoli, G., Rinaldi, L., Monteiro, M. F. M., Calado, A. M. C., Ramos, R. A. N., Meira-Santos, P. O., & Alves, L. C. (2015). A comparison of Mini-FLOTAC and FLOTAC with classic methods to diagnosing intestinal parasites of dogs from Brazil. *Parasitology Research*, *114*, 3529-3533.
- Livia, K. G., Reyes, R., Ramos, V. A., González, E. B., Carrillo, N. M., Ponce, E. R., & Foronda, P. (2023). Metastrongyloid Infection with *Aelurostrongylus abstrusus*, *Troglostrongylus brevior*, *Oslerus rostratus* and *Angiostrongylus chabaudi* in Feral Cats from the Canary Islands (Spain). *Animals*, *13*, 1-16.
- Marchiori, E., Bono, L., Voltan, L., Dotto, G., Tessarin, C., & Marcer, F. (2024). Gastrointestinal Parasites in Non-Human Primates in Zoological Gardens in Northern Italy. *Animals*, *14*, 1-12.
- Mcglade, T. R., Robertson, I. D., Elliot, A. D., Read, C., & Thompson, R. C. (2003). Gastrointestinal parasites of domestic cats in Perth, Western Australia. *Veterinary Parasitology*, *28*, 251-262.
- Monteiro, M. F., Ramos, R. A., Calado, A. M., Lima, V. F., Ramos, I. C., & Tenório, R. F., Faustino, M. A., & Alves, L. C. (2016). Gastrointestinal parasites of cats in Brazil: frequency and zoonotic risk. *Revista Brasileira de Parasitologia Veterinária*, *25*, 254-257.
- Moreira, A. D. S., Baptista, C. T., Brasil, C. L., Valente, J. S. S., Bruhn, F. R. P., & Pereira, D. I. B. (2018). Risk factors and infection due to *Cryptosporidium* spp. in dogs and cats in southern Rio Grande do Sul. *Revista Brasileira de Parasitologia Veterinária*, *27*, 113-118.
- Morelli, S., Traversa, D., Diakou, A., Colombo, M., Russi, I., Mestek, A., Chandrashekar, R., Beall, M., Paoletti, B., & Iorio, R. (2022). A comparison of copromicroscopic and molecular methods for the diagnosis of cat *Aelurostrongylosis*. *Animals*, *12*, 1-10.
- Napoli, E., Pugliese, M., Basile, A., Passantino, A., & Brianti, E. (2023). *Clinical*, radiological, and echocardiographic findings in cats infected by *Aelurostrongylus abstrusus*. *Pathogens*, *12*, 1-13.
- Nonnis, F., Tamponi, C., Tosciri, G., Manconi, M., Pudda, F., Cabras, P., Dessi, G., Scala, A., & Varcasia, A. (2023). Cardio-pulmonary nematodes of the red fox (*Vulpes vulpes*) of Sardinia, Italy. *Parasitology Research*, *122*, 1685-1688.
- Pereira, P. F., Barbosa, A. D. S., Moura, A. P. P., Vasconcellos, M. L., Uchôa, C. M. A., Bastos, O. M. P., & Amendoeira, M. R. R. (2017). Gastrointestinal parasites in stray and shelter cats in the municipality of Rio de Janeiro, Brazil. *Revista Brasileira de Parasitologia Veterinária*, *26*, 383-388.
- Pfukenyi, D. M., Chipunga, S. L., Dinginya, L., & Matenga, E. (2010). A survey of pet ownership, awareness and public knowledge of pet zoonoses with particular reference to roundworms and hookworms in Harare, Zimbabwe. *Tropical Animal Health and Production*, *42*, 247-252.

- Sauda, F., Malandrucchio, L., Liberato, C., & Perrucci, S (2019). *Gastrointestinal parasites in shelter cats of central Italy. Veterinary Parasitology Regional Studies Report*, 18, 1-5.
- Silva, Y. H., Campos, D. R., Lima, G. A. C., Quintal, J. P., Guimarães, B. G., Rêgo, G. M. M., Avelar, B. R., Intriери, J. M., Correia, T. R., & Scott, F. B (2023). Prevalence of gastrointestinal parasites in domestic cats (*Felis catus*) diagnosed by different coproparasitological techniques in the municipality of Seropédica, Rio de Janeiro. *Brazilian Journal Veterinary Parasitology*, 32, 1-9.
- Simpson, J. W. (1998). Diet and large intestinal disease in dogs and cats. *The Journal of Nutrition*, 128(12 Suppl), 2717-2722.
- Stalliviere, F. M., Bellato, V., Souza, A. P., Sartor, A. A., Moura, A. B., & Rosa, L. D. (2009). Ectoparasites and intestinal helminths in *Felis catus domesticus* from Lages city, SC, Brazil and social-economical and cultural aspects of owners of family pets. *Revista Brasileira de Parasitologia Veterinária*, 18, 26-30.
- Stepanović, P., Despotović, D., Dimitrijević, S., & Ilić, T. (2020). Clinical-parasitological screening for respiratory capillariasis in cats in urban environments. *Helminthology*, 19, 322-334.
- Symeonidou, I., Gelasakis, A. I., Arsenopoulos, K., Angelou, A., Beugnet, F., & Papadopoulos, E. (2018). Feline gastrointestinal parasitism in Greece: emergent zoonotic species and associated risk factors. *Parasites & Vectors*, 4, 1-13.
- Taylor, M. A., Coop, R. L., & Wall, R. L. (2010). *Parasitologia Veterinária*. ed.3. Guanabara Koogan.
- Traversa, D., Di Cesare, A., & Conboy, G. (2010). Canine and feline cardiopulmonary parasitic nematodes in Europe: emerging and underestimated. *Parasites & Vectors*, 3, 1-22.
- Traversa, D., & Guglielmini, C. (2008). Feline aelurostrongylosis and canine angiostrongylosis: a challenging diagnosis for two emerging verminous pneumonia infections. *Veterinary Parasitology*, 157, 163-174.
- Vincy, P., & Tresamol, P. V (2023). Prevalence of gastro-intestinal and haemoparasitic infections among domestic cats of Kerala. *Journal of Parasitic Diseases*, 47, 562-565.
- Vismarra, A., Schnyder, M., Strube, C., Kramer, L., Colombo, L., & Genchi, M (2023). Diagnostic challenges for *Aelurostrongylus abstrusus* infection in cats from endemic areas in Italy. *Parasites & Vectors*, 16, 1-6.
- Willesen, J. (2021). *The Baermann test for suspected Angiostrongylosis*. Bayer Animal Health GmbH, 51368 Leverkusen, Germany, 2009. www.animalhealth.bayerhealthcare.com/http://lungenwürmer.bayer.de/static/media/BaermannFlyer.pdf.
- Yang, Y., & Liang, H. (2015). Prevalence and risk factors of intestinal parasites in cats from China. *BioMed Research International*, 2015, 1-6.
- Yildirim, I., Ozturk, B., Solmaz, D. E., Baydar, E., & Aydogdu, U. (2023). Determination of *Aelurostrongylus abstrusus* prevalence and risk factors in cats from Balikesir. *Turkiye Parazitoloji Dergisi*, 47, 78-82.

Received January 8, 2026.

Accepted February 11, 2026.